


Meisoindigo

Chemical Properties

CAS No. :	97207-47-1
Formula:	C ₁₇ H ₁₂ N ₂ O ₂
Molecular Weight:	276.29
Storage:	Powder: -20°C for 3 years In solvent: -80°C for 1 year Actual storage temperature shall be subject to the COA.



Biological Description

Description	Meisoindigo (Natura- α) is a derivative of indigo natural, might induces apoptosis and myeloid differentiation of acute myeloid leukemia (AML).
Targets(IC50)	Apoptosis
Kinase Assay	GSK3 Scintillation Proximity Assay: The competition experiments are carried out in duplicate with 10 concentrations of the inhibitor in clear-bottomed microtiter plates. The biotinylated peptide substrate, biotin-AAEELDSRAGS(PO ₃ H ₂)PQL, is added at a final concentration of 2 μ M in an assay buffer containing 6 milliunits of recombinant human GSK3 (equal mix of both α and β), 12 mM MOPS, pH 7.0, 0.3 mM EDTA, 0.01% β -mercaptoethanol, 0.004% Brij 35, 0.5% glycerol, and 0.5 μ g of bovine serum albumin/25 μ l and preincubated for 10-15 min. The reaction is initiated by the addition of 0.04 μ Ci of [γ - ³³ P]ATP and unlabeled ATP in 50 mM Mg(Ac) ₂ to a final concentration of 1 μ M ATP and assay volume of 25 μ l. Blank controls without peptide substrate are used. After incubation for 20 min at room temperature, each reaction is terminated by the addition of 25 μ l of stop solution containing 5 mM EDTA, 50 μ M ATP, 0.1% Triton X-100, and 0.25 mg of streptavidin-coated SPA beads corresponding to ~ 35 pmol of binding capacity. After 6 h the radioactivity is determined in a liquid scintillation counter. Inhibition curves are analyzed by non-linear regression using GraphPad Prism.

Solubility Information

Solubility	DMSO: 18.33 mg/mL (66.34 mM),Sonication is recommended. (< 1 mg/ml refers to the product slightly soluble or insoluble)
In vivo Formulation	10% DMSO+40% PEG300+5% Tween 80+45% Saline: 2 mg/mL (7.24 mM),Sonication is recommended. <i>Please add the solvents sequentially, clarifying the solution as much as possible before adding the next one. Dissolve by heating and/or sonication if necessary. Working solution is recommended to be prepared and used immediately. The formulation provided above is for reference purposes only. In vivo formulations may vary and should be modified based on specific experimental conditions.</i>

Preparing Stock Solutions

	1mg	5mg	10mg
1 mM	3.6194 mL	18.0969 mL	36.1939 mL
5 mM	0.7239 mL	3.6194 mL	7.2388 mL
10 mM	0.3619 mL	1.8097 mL	3.6194 mL
50 mM	0.0724 mL	0.3619 mL	0.7239 mL

Please select the appropriate solvent to prepare the stock solution, according to the solubility of the product in different solvents. Please use it as soon as possible.

Note: The dilution table applies only to solid products. For liquid products, please calculate the stock solution based on the stated concentration and/or density.

Reference

Lee CC, et al. Leuk Lymphoma. 2010 May;51(5):897-905.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481