

HGFR/c-Met Protein, Human, Recombinant (His), PE-Labeled

General Information

Synonyms:	c-Met;RCCP2;HGFR;DFNB97;DA11;AUTS9
Protein Construction:	A DNA sequence encoding the Human MET (NP_000236.2) (Met1-Thr932) was expressed with a polyhistidine tag at the C-terminus. The protein is site-specifically labeled with PE (Excitation = 495-566 nm, Emission Max. = 575 nm).
Species:	Human
Expression Host:	HEK293 Cells
Accession:	P08581-1
Molecular Weight:	105.1 kDa (predicted)

QC Testing

Biological Activity:	<p>1. Flow cytometric analysis of anti-c-MET CAR expression. 293 cells were lentivirally transduced with anti-c-MET CAR. Flow cytometric analysis was performed with a negative control protein and HGFR/c-Met Protein, Human, Recombinant (His), PE-Labeled (Cat#TMPY-07027), respectively. Non-transduced 293 cells were used as a control (left).
</p> <p>2. Binding activity of PE-conjugated c-MET protein to PBMC cells. PBMC cells were stained with anti-CD3 antibody and HGFR/c-Met Protein, Human, Recombinant (His), PE-Labeled (Cat#TMPY-07027) and detected by flow cytometry. PBMC cells stained with anti-CD3 antibody were used as a control (left).</p>
Endotoxin:	< 1.0 EU/μg of the protein as determined by the LAL method.
Formulation:	This product is Lyophilized from sterile PBS, 0.2% BSA, pH 7.4. Please contact us for any concerns or special requirements. Please refer to the specific buffer information in the hardcopy of datasheet or the lot-specific COA.

Preparation and Storage

Reconstitution:	Please refer to the lot-specific COA.
Stability & Storage:	Twelve months from date of receipt at -20°C to -70°C in lyophilized form and 3 months at -70°C under sterile conditions after reconstitution. Protect from prolonged exposure to light and avoid repeated freeze-thaw cycles. <small>Actual storage temperature shall be subject to the COA.</small>
Shipping:	In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

Protein Background

Hepatocyte growth factor receptor (HGFR), also known as c-Met or mesenchymal-epithelial transition factor (MET), is a receptor tyrosine kinase (RTK) that is overexpressed and/or mutated in a variety of malignancies. HGFR protein

is produced as a single-chain precursor, and HGF is the only known ligand. Normal HGF/HGFR signaling is essential for embryonic development, tissue repair, or wound healing, whereas aberrantly active HGFR has been strongly implicated in tumorigenesis, particularly in the development of invasive and metastatic phenotypes. HGFR protein is a multifaceted regulator of growth, motility, and invasion, and is normally expressed by cells of epithelial origin. Preclinical studies suggest that targeting aberrant HGFR signaling could be an attractive therapy in cancer. Cancer Immunotherapy Immune Checkpoint Immunotherapy Targeted Therapy

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel: 781-999-4286 E_mail: info@targetmol.com Address: 34 Washington Street, Wellesley Hills, MA 02481