


C188-9

Chemical Properties

CAS No. :	432001-19-9
Formula:	C ₂₇ H ₂₁ NO ₅
Molecular Weight:	471.52
Storage:	Store under nitrogen Powder: -20°C for 3 years In solvent: -80°C for 1 year <i>Actual storage temperature shall be subject to the COA.</i>



Biological Description

Description	C188-9 (TTI-101) is a Stat3 inhibitor with an IC ₅₀ of 4-7 μM.
Targets(IC ₅₀)	Apoptosis,STAT
In vitro	For apoptosis studies, AML cell lines and primary samples are treated for 24 hours with C188-9, then apoptotic cells are quantified by FACS analysis for annexin V-labeled cells. The EC ₅₀ s for apoptosis induction are quite variable, ranging from 6 μM to over 50 μM
In vivo	In an analysis of approximately 13,528 discernible genes, C188 influences the expression of 37 gene transcripts (17 downregulated and 20 upregulated, false discovery rate (fdr)<0.01, fold change≥1.5), including 7 known targets of STAT3. Conversely, C188-9 exhibits a broader impact, modifying the expression of 384 genes related to oncogenesis (95 down- and 289 up-regulated), 76 of which are regulated by STAT3 (38 down-regulated and 38 up-regulated). Notably, C188-9 treatment downregulates 24 out of 38 genes (63%) previously identified as upregulated by STAT3. Furthermore, C188-9 downregulates an additional 10 genes (fdr <0.01, fold change≥1.5) known to be upregulated by STAT1, suggesting that 83.3% (40 out of 48) of the genes downregulated by C188-9 are positively regulated by STAT1. This includes sixteen genes co-regulated by both STAT3 and STAT1. The findings suggest that C188-9's modulation of gene transcript levels in head and neck squamous cell carcinoma (HNSCC) tumors may be attributed to its dual influence on STAT3 and STAT1 pathways.

Solubility Information

Solubility	DMSO: 262.5 mg/mL (556.71 mM),Sonication is recommended. (< 1 mg/ml refers to the product slightly soluble or insoluble)
In vivo Formulation	10% DMSO+40% PEG300+5% Tween 80+45% Saline: 6 mg/mL (12.72 mM),Solution. <i>Please add the solvents sequentially, clarifying the solution as much as possible before adding the next one. Dissolve by heating and/or sonication if necessary. Working solution is recommended to be prepared and used immediately. The formulation provided above is for reference purposes only. In vivo formulations may vary and should be modified based on specific experimental conditions.</i>

Preparing Stock Solutions

	1mg	5mg	10mg
1 mM	2.1208 mL	10.604 mL	21.208 mL
5 mM	0.4242 mL	2.1208 mL	4.2416 mL
10 mM	0.2121 mL	1.0604 mL	2.1208 mL
50 mM	0.0424 mL	0.2121 mL	0.4242 mL

Please select the appropriate solvent to prepare the stock solution, according to the solubility of the product in different solvents. Please use it as soon as possible.

Note: The dilution table applies only to solid products. For liquid products, please calculate the stock solution based on the stated concentration and/or density.

Reference

edell MS, et al. Stat3 signaling in acute myeloid leukemia: ligand-dependent and -independent activation and induction of apoptosis by a novel small-molecule Stat3 inhibitor. *Blood*. 2011 May 26;117(21):5701-9.

Gu H, Chen C, Hao X, et al. MDH1-mediated malate-aspartate NADH shuttle maintains the activity levels of fetal liver hematopoietic stem cells. *Blood*. 2020, 136(5): 553-571.

He, Qiu-Rui, et al. The natural product trienomycin A is a STAT3 pathway inhibitor that exhibits potent in vitro and in vivo efficacy against pancreatic cancer. *British Journal of Pharmacology*. 178.12 (2021): 2496-2515.

Bharadwaj U, et al. Small-molecule inhibition of STAT3 in radioresistant head and neck squamous cell carcinoma. *Oncotarget*. 2016 May 3;7(18):26307-30.

He Q R, Tang J J, Chen Z F, et al. Natural Product Trienomycin A is a STAT3 pathway inhibitor that exhibits potent in vitro and in vivo efficacy against pancreatic cancer[J]. *Authorea Preprints*. 2020

Guan X, Yang J, Wang W, et al. Dual inhibition of MYC and SLC39A10 by a novel natural product STAT3 inhibitor derived from *Chaetomium globosum* suppresses tumor growth and metastasis in gastric cancer. *Pharmacological Research*. 2023: 106703.

Gu H, Chen C, Hao X, et al. MDH1-mediated malate-aspartate NADH shuttle maintains the activity levels of fetal liver hematopoietic stem cells[J]. *Blood*. 2021, 137(11): 1478-1490.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel: 781-999-4286 E_mail: info@targetmol.com Address: 34 Washington Street, Wellesley Hills, MA 02481