

CCL15 Protein, Human, Recombinant

General Information

Synonyms:	Small-inducible cytokine A15;Mrp-2b;NCC3;NCC-3;Leukotactin-1 (LKN-1);Macrophage inflammatory protein 5 (MIP-5);C-C motif chemokine 15;SCYA15;CCL15;MIP5;MIP-1 delta; Chemokine CC-2 (HCC-2)
Protein Construction:	22-113 aa
Species:	Human
Expression Host:	E. coli
Accession:	Q16663
Molecular Weight:	10.2 kDa (Predicted)
AA Sequence:	QFINDAETELMMSKLPLENPVVLSNFHFAADCCTSYISQIPCSLMKSYFETSSECSKPGVIFLTKKGRQVCAKP SGPGVQDCMKKLPYSI

QC Testing

Biological Activity:	Fully biologically active when compared to standard. The biological activity determined by a chemotaxis bioassay using human T-lymphocytes is in a concentration range of 1.0-10 ng/ml.
Purity:	>97% as determined by SDS-PAGE.
Endotoxin:	< 1.0 EU/μg of the protein as determined by the LAL method.
Formulation:	Lyophilized from a 0.2 μm filtered PBS, pH 7.4

Preparation and Storage

Reconstitution:

Reconstitute the lyophilized protein in sterile deionized water. The product concentration should not be less than 100 μg/ml. Before opening, centrifuge the tube to collect powder at the bottom. After adding the reconstitution buffer, avoid vortexing or pipetting for mixing.

Stability & Storage:

Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Actual storage temperature shall be subject to the COA.

Shipping:

In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481