

## Anti-4-1BB/CD137/TNFRSF9 Antibody-FITC (9J469)

### Product Details

Ig Type:	Rabbit IgG
Reactivity:	Human
Conjugation:	FITC
Clone:	9J469
Purification:	Protein A

### Applications

Verified Activity:	Flow cytometric analysis of Human CD137 expression on PHA-activated human whole blood lymphocytes. Cells were stained with FITC-conjugated anti-Human CD137. The fluorescence histograms were derived from gated events with the forward and side light-scatter characteristics of viable lymphocytes.
Application:	FCM
Recommended	10 µl/Test, 0.1 mg/ml

### Properties

Stability & Storage:	Store at 2°C-8°C for 12 months, do not freeze. Keep away from direct sunlight.
Shipping:	Shipping with blue ice.

### Antigen Details

Immunogen:	Recombinant Protein: Human CD137 Protein (TMPY-01743)
Antigen Species:	Human
Synonyms:	A930040I11Rik;CDw137;AI325004;tumor necrosis factor receptor superfamily, member 9;ILA;Ly63;4-1BB;Cd137;AA408498
Biology Area:	Cancer Drug Targets

### Research Background

CD137 (also known as 4-1BB) is a surface co-stimulatory glycoprotein originally described as present on activated T lymphocytes, which belongs to the tumor necrosis factor (TNF) receptor superfamily. It is expressed mainly on activated CD4+ and CD8+ T cells, and binds to a high-affinity ligand (4-1BBL) expressed on several antigen-presenting cells such as macrophages and activated B cells. Upon ligand binding, 4-1BB is associated with the tumor necrosis factor receptor-associated factors (TRAFs), the adaptor protein which mediates downstream signaling events including the activation of NF- $\kappa$ B and cytokine production. 4-1BB signaling either by binding to 4-1BBL or by antibody ligation delivers signals for T-cell activation and growth, as well as monocyte proliferation and B-cell survival, and plays an important role in the amplification of T cell-mediated immune responses. In addition, CD137 and CD137L are expressed in different human primary tumor tissues, suggesting that they may influence the progression of tumors. Crosslinking of CD137 on activated T cells has shown promise in enhancing anti-tumor immune responses in murine models, and agonistic anti-CD137 antibodies are currently being tested in phase I clinical trials. Soluble forms of CD137 (sCD137) are generated by differential splicing. sCD137 can bind to CD137 ligand to antagonize the costimulatory activities of the membrane-bound CD137 and reduce T cell proliferation and IL-2 secretion. Cancer Immunotherapy Co-stimulatory Immune Checkpoint Targets Immune

## A DRUG SCREENING EXPERT

---

Checkpoint Detection: AntibodiesImmune Checkpoint Detection: ELISA AntibodiesImmune Checkpoint  
ProteinsImmune Checkpoint TargetsImmunotherapyTargeted Therapy

**Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins**

This product is for Research Use Only· Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E\_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481