

Anti-Aminopeptidase N/CD13 Antibody (8T921)

Product Details

Ig Type:	Rabbit IgG
Reactivity:	Human
Conjugation:	Unconjugated
Clone:	8T921
Purification:	Protein A

Applications

Verified Activity:	<ol style="list-style-type: none">1. Immunochemical staining of human ANPEP in human renal carcinoma with rabbit monoclonal antibody (1:200, formalin-fixed paraffin embedded sections).2. Immunochemical staining of human ANPEP in human liver with rabbit monoclonal antibody (1:200, formalin-fixed paraffin embedded sections).3. Anti-ANPEP rabbit monoclonal antibody at 1:500 dilution.<ul style="list-style-type: none">-Lane A: THP1 Whole Cell Lysate.-Lane B: U937 Whole Cell lysate. <ul style="list-style-type: none">-Lysates/proteins at 30 µg per lane.-Secondary<ul style="list-style-type: none">-Goat Anti-Rabbit IgG H&L (Dylight800) at 1/10000 dilution.-Developed using the Odyssey technique.-Performed under reducing conditions.-Predicted band size:110 kDa.-Observed band size:140 kDa
Application:	ELISA,IHC-P,WB
Recommended	WB: 1:500-1:2000; ELISA: 1:5000-1:10000; IHC-P: 1:100-1:500

Properties

Stability & Storage:	Store at 2°C-8°C for 1 month. Store at -20°C or -80°C for 12 months. Avoid repeated freeze-thaw cycles. Preservative-Free.
Shipping:	Shipping with blue ice.

Antigen Details

Immunogen:	Recombinant Protein: Human CD13 / ANPEP protein (TMPY-00665)
Antigen Species:	Human
Synonyms:	P150;alanyl (membrane) aminopeptidase;AP-M;AP-N;Cd13;Apn

Research Background

Aminopeptidase N (ANPEP or APN), also known as CD13, is a cell-surface metalloprotease located in the small-intestinal and renal microvillar membrane, as well as other plasma membranes. It belongs to the peptidase M1 family. CD13 plays a role in the final digestion of peptides generated from hydrolysis of proteins by gastric and pancreatic proteases and is involved in the metabolism of regulatory peptides by diverse cell types. CD13/APN is a potent regulator of angiogenesis which is essential for tumor invasion and metastasis, and its transcription in

A DRUG SCREENING EXPERT

activated endothelial cells is induced by angiogenic growth factors via the RAS/MAPK pathway. In addition, this enzyme has been shown to participate in antigen processing and presentation, and accordingly, defects in this gene appear to be a cause of various types of leukemia or lymphoma and carcinomas.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481