

Anti-CD25/IL2R alpha Antibody-PerCP (4X703)

Product Details

| | |
|---------------|------------|
| Ig Type: | Mouse IgG1 |
| Reactivity: | Human |
| Conjugation: | PerCP |
| Clone: | 4X703 |
| Purification: | Protein A |

Applications

| | |
|--------------------|--|
| Verified Activity: | Flow cytometric analysis of CD25/IL2R alpha expression on stimulated human peripheral blood lymphocytes. Human peripheral blood mononuclear cells were stimulated for 3 days with Phytohemagglutinin, then stained with PerCP-conjugated anti-Human CD25/IL2R alpha. The fluorescence histograms were derived from events with the forward and side light-scatter characteristics of viable lymphocytes. |
| Application: | FCM |
| Recommended | 5 μ L/Test, 0.1 mg/mL |

Properties

| | |
|----------------------|--|
| Stability & Storage: | Store at 2°C-8°C for 12 months, do not freeze. Keep away from direct sunlight. |
| Shipping: | Shipping with blue ice. |

Antigen Details

| | |
|------------------|---|
| Immunogen: | Recombinant Protein: Human CD25/IL2R alpha Protein (TMPY-01103) |
| Antigen Species: | Human |
| Synonyms: | CD25/IL2R α ;interleukin 2 receptor, α ;interleukin 2 receptor, alpha |
| Biology Area: | Neuroinflammation, Cancer Drug Targets |

Research Background

CD25 (alpha-chain of the IL-2 receptor, or IL2RA), is a type I transmembrane glycoprotein with a signal peptide, an extracellular region, a transmembrane region, and a cytoplasmic domain. IL2RA is expressed on activated T cells and regulatory T cells and is capable of binding IL2 with low affinity by itself. However, a ligand-induced high-affinity heterotrimeric receptor complex is produced when IL2RA is associated non-covalently with the IL2 receptor beta and gamma chain, and subsequently initiates the intracellular signal pathways such as MAPK or JAK/STAT. On dendritic cells (DC), CD25 has been previously regarded as an activation marker, while both murine and human DC can express CD25, they do not express the beta-chain of the IL-2 receptor, which is indispensable for the execution of IL-2 signaling. The IL2RA (CD25) gene is a substantial component of the high-affinity receptor molecule highly expressed by activated T lymphocytes. Recently, a piece of strong evidence was obtained for the involvement of IL-2RA in conferring susceptibility to type 1 diabetes (T1D). Cancer growth and development are associated with the stimulation of the innate immune system, including enhanced interleukin 2 receptor (IL-2R) expression in immune cells and its shedding into the circulation in a soluble form of SIL-2Ralpha. In most hematological malignancies, including different types of leukemias and lymphomas, SIL-2Ralpha is released directly from the surface of

A DRUG SCREENING EXPERT

neoplastic cells thus reflecting the tumor bulk, turnover, and activity. Several studies have proved that not only lymphoid cancer cells but also some non-lymphoid cancer cells, express IL-2R on their surface. They include malignant melanoma and carcinomas of the kidney, head and neck, esophagus, and lung. Thus, sIL-2R α is elevated in most proliferative disturbances of the hematopoietic system and many solid tumors. Cancer Immunotherapy Immune Checkpoint Immunotherapy Targeted Therapy

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel: 781-999-4286 E_mail: info@targetmol.com Address: 34 Washington Street, Wellesley Hills, MA 02481