

Anti-CD45 Antibody (5N468)

Product Details

Ig Type:	Mouse IgG1
Reactivity:	Human
Conjugation:	Unconjugated
Clone:	5N468
Purification:	Protein A

Applications

1. Immunochemical staining of human PTPRC in human lymphoma with mouse monoclonal antibody at 1:5000 dilution, formalin-fixed paraffin embedded sections.
2. Immunochemical staining of human PTPRC in human spleen with mouse monoclonal antibody at 1:5000 dilution, formalin-fixed paraffin embedded sections.
3. Anti-PTPRC mouse monoclonal antibody at 1:500 dilution.

Verified Activity:	-Lane A: Jurkat Whole Cell Lysate.
	-Lane B: Daudi Whole Cell lysate.
	-Lysates/proteins at 30 µg per lane.
	-Secondary
	-Goat Anti-Mouse IgG (H+L)/HRP at 1/10000 dilution.
Application:	IHC-P,WB
Recommended	WB: 1:500-1:2000; IHC-P: 1:1000-1:10000

Properties

Stability & Storage:	Store at 2°C-8°C for 1 month. Store at -20°C or -80°C for 12 months. Avoid repeated freeze-thaw cycles. Preservative-Free.
Shipping:	Shipping with blue ice.

Antigen Details

Immunogen:	A synthetic peptide: center region of the Human CD45/PTPRC
Antigen Species:	Human
Synonyms:	LCA;GP180;L-CA;CD45;B220;CD45R;protein tyrosine phosphatase, receptor type, C;T200;LY5
Biology Area:	Phosphatases and Regulators

Research Background

The cluster of differentiation (CD) system is commonly used as cell markers in Immunophenotyping. Different kinds of cells in the immune system can be identified through the surface CD molecules associating with the immune function of the cell. There are more than 320 CD unique clusters and subclusters have been identified. Some of the CD molecules serve as receptors or ligands important to the cell through initiating a signal cascade which then alter the behavior of the cell. Some CD proteins do not take part in cell signal process but have other functions such as cell

adhesion. Protein tyrosine phosphatase, receptor type C (CD45), also known as PTPRC is a member of the protein tyrosine phosphatase (PTP) family which is known for its function to serve as signaling molecules and to regulate a variety of cellular processes such as cell proliferation, differentiation, mitotic cycle and oncogenic transformation. CD45 is found expression specifically in hemotopietic cells. CD45 consists of an extracellular domain, a single transmembrane segment and two tandem intracytoplasmic catalytic domains. It serves as an essential regulator of T-cell and B-cell antigen receptor signaling through either direct interaction with components of the antigen receptor complexes or by activating various Src family kinases required for the antigen receptor signaling and it also can suppress JAK kinases.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only· Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481