

## Anti-CD50 Antibody-PE (5P32)

## Product Details

|               |             |
|---------------|-------------|
| Ig Type:      | Mouse IgG2a |
| Reactivity:   | Human       |
| Conjugation:  | PE          |
| Clone:        | 5P32        |
| Purification: | Protein A   |

## Applications

|                    |   |
|--------------------|---|
| Verified Activity: | Profile of peripheral blood lymphocytes analyzed by flow cytometry. The fluorescence histograms were derived from events with the forward and side light-scatter characteristics of viable lymphocytes. |
| Application:       | FCM   |
| Recommended        | 5 µl/Test, 0.1 mg/ml  |

## Properties

|                      |  |
|----------------------|--|
| Stability & Storage: | Store at 2°C-8°C for 12 months, do not freeze. Keep away from direct sunlight. Sodium azide is toxic to cells and should be disposed of properly. Flush with large volumes of water during disposal. |
| Shipping:            | Shipping with blue ice.  |

## Antigen Details

|                  |  |
|------------------|--|
| Immunogen:       | Recombinant Protein: Human ICAM3 / CD50 protein (TMPY-01430) |
| Antigen Species: | Human  |
| Synonyms:        | CD50;CDW50;ICAM-R;ICAM-3;intercellular adhesion molecule 3   |

## Research Background

The protein ICAM-3, also known as CD50, is a member of the intercellular adhesion molecule (ICAM) family consisting of three members. It is a DC-SIGN ligand that is constitutively expressed on resting leukocytes and is thus an important molecule for the first immune response. ICAM-3 comprises five immunoglobulin-like domains and binds LFA-1 through its two N-terminal domains. It functions not only as an adhesion molecule but also as a potent signaling molecule. ICAM-3 binds to LFA-1 on antigen-presenting cells (APC) stabilizing the T cell-APC interaction, facilitating signaling through the CD3/TCR complex. However, recent evidence using cultured and transformed T cells suggests ICAM-3 may also function in signaling. It has been reported that the CD50 molecule can play a role in developing functionally mature T lymphocytes and its expression increases during the maturation process of T lymphocytes. Also, the interactions of ICAM-3 and LFA-1 facilitate HIV-1- induced virological synapse formation between T cells. ICAM-3 is associated with an increase in cellular radio-resistance and cancer cell proliferation. It could be considered as a candidate for anti-cancer drug development and as a cancer diagnostic marker.

**Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins**

This product is for Research Use Only. Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E\_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481