

## Anti-IFNGR1 Antibody (3T547)

## Product Details

Ig Type:	Rabbit IgG
Reactivity:	Human
Conjugation:	Unconjugated
Clone:	3T547
Purification:	Protein A

## Applications

	Anti-IFNGR1 rabbit monoclonal antibody at 1:500 dilution. -Lane A: MCF7 Whole Cell lysate. -Lysates/proteins at 30 µg per lane. -Secondary
Verified Activity:	-Goat Anti-Rabbit IgG H&L (Dylight800) at 1/10000 dilution. -Developed using the Odyssey technique. -Performed under reducing conditions. -Predicted band size:54 kDa. -Observed band size:54 kDa
Application:	ELISA,WB
Recommended	WB: 1:500-1:1000; ELISA: 1:25000-1:50000

## Properties

Stability & Storage:	Store at 2°C-8°C for 1 month. Store at -20°C or -80°C for 12 months. Avoid repeated freeze-thaw cycles. Preservative-Free.
Shipping:	Shipping with blue ice.

## Antigen Details

Immunogen:	Recombinant Protein: Human CD119 protein (TMPY-01151)
Antigen Species:	Human
Synonyms:	interferon gamma receptor 1;interferon γ receptor 1;CD119;IMD27A;IMD27B;IFNGR

## Research Background

The cluster of differentiation (CD) system is commonly used as cell markers in Immunophenotyping. Different kinds of cells in the immune system can be identified through the surface CD molecules associating with the immune function of the cell. There are more than 320 CD unique clusters and subclusters have been identified. Some of the CD molecules serve as receptors or ligands important to the cell through initiating a signal cascade which then alters the behavior of the cell. Some CD proteins do not take part in the cell signal process but have other functions such as cell adhesion. CD119 (cluster of differentiation 119), also known as IFNGR1 (interferon-gamma receptor 1), is part of the heterodimeric gamma interferon receptor which consists of IFNGR1 (CD119) and IFNGR2. The IFNGR1 gene encodes the ligand-binding chain (alpha) of the interferon receptor while the IFNGR gene encodes the non-ligand binding partner. The ability of the interferon-γ was achieved through binding to the interferon receptor CD119. After binding, the products of activated T-lymphocytes interferon-γ exerts antiviral activity, growth inhibitory effect, and

several immune- regulatory activities on a variety of cell types.

**Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins**

This product is for Research Use Only· Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E\_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481