

Anti-SMAC Antibody (5G346)

Product Details

Ig Type:	Rabbit IgG
Reactivity:	Human
Conjugation:	Unconjugated
Clone:	5G346
Purification:	Protein A

Applications

Diablo was immunoprecipitated using:

- Lane A:0.5 mg 293T Whole Cell Lysate.
- Lane B:0.5 mg MCF-7 Whole Cell Lysate.
- Lane C:0.5 mg HepG2 Whole Cell Lysate.

-2 µL anti-Diablo rabbit monoclonal antibody and 15 µl of 50 % Protein G agarose.

-Primary antibody:

Verified Activity: -Anti-Diablo rabbit monoclonal antibody, at 1:330 dilution.

-Secondary antibody:

- Dylight 800-labeled antibody to rabbit IgG (H+L), at 1:5000 dilution.
- Developed using the odyssey technique.
- Performed under reducing conditions.
- Predicted band size: 19 kDa.
- Observed band size: 19 kDa

Application: IP

Recommended IP: 0.5-2 µL/mg of lysate

Properties

Stability & Storage: Store at 2°C-8°C for 1 month. Store at -20°C or -80°C for 12 months. Avoid repeated freeze-thaw cycles. Preservative-Free.

Shipping: Shipping with blue ice.

Antigen Details

Immunogen: Recombinant Protein: Human SMAC / Diablo Protein (TMPY-00924)

Antigen Species: Human

Synonyms: SMAC;DFNA64;diablo, IAP-binding mitochondrial protein

Research Background

Apoptosis is an essential processes required for normal development and homeostasis of all metazoan organisms. Second Mitochondria-Derived Activator of Caspases (Smac) or Direct IAP Binding Protein with low isoelectric point, pI (Diablo) is a proapoptogenic mitochondrial protein that is released to the cytosol in response to diverse apoptotic stimuli, including commonly used chemotherapeutic drugs. The current knowledge about structure and function of Smac/Diablo during programmed cell death, both in mitochondrial and receptor pathways are presented. It has been shown that Diablo mainly interacts with IAPs in the cytochrome c/Apaf-1/caspase-9 pathway, and promotes

apoptosis. Diablo is released from the mitochondria into the cytosol occurring downstream of cytochrome c release in response to apoptotic stimuli such as irradiation, DNA damage or cytotoxic drugs. In the cytosol, Smac/Diablo interacts and antagonizes inhibitors of apoptosis proteins (IAPs), thus allowing the activation of caspases and apoptosis. This activity has prompted the synthesis of peptidomimetics that could potentially be used in cancer therapy. The role of Smac/DIABLO in colorectal carcinogenesis is ill defined. Data continues to accumulate to suggest that decreased levels of Smac/DIABLO may be important in chemoradiation-resistance to apoptosis in advanced colon cancer.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only· Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481