

## Anti-FLT3 Antibody-FITC (2N391)

## Product Details

Ig Type:	Mouse IgG1
Reactivity:	Human
Conjugation:	FITC
Clone:	2N391
Purification:	Protein A

## Applications

Verified Activity:	Flow cytometric analysis of anti-human CD135 on human whole blood monocytes. The fluorescence histograms were derived from events with the forward and side light-scatter characteristics of viable monocytes.
Application:	FCM
Recommended	10 µl/Test, 0.1 mg/ml

## Properties

Stability & Storage:	Store at 2°C-8°C for 12 months, do not freeze. Keep away from direct sunlight. Sodium azide is toxic to cells and should be disposed of properly. Flush with large volumes of water during disposal.
Shipping:	Shipping with blue ice.

## Antigen Details

Immunogen:	Recombinant Protein: Human FLT3 / CD135 / FLK2 protein (TMPY-01247)
Antigen Species:	Human
Synonyms:	FLK-2;fms-related tyrosine kinase 3;FLK2;STK1;CD135
Biology Area:	Cancer Drug Targets, Receptor Tyrosine Kinases (RTKs)

## Research Background

The cluster of differentiation (CD) system is commonly used as cell markers in Immunophenotyping. Different kinds of cells in the immune system can be identified through the surface CD molecules which associating with the immune function of the cell. There are more than 320 CD unique clusters and subclusters have been identified. Some of the CD molecules serve as receptors or ligands important to the cell through initiating a signal cascade which then alter the behavior of the cell. Some CD proteins do not take part in cell signal process but have other functions such as cell adhesion. CD135, also known as FLT-3, FLK-2, is a member of the CD system. CD135 is an important cell surface marker recognized by specific sets of antibodies to identify the types of hematopoietic (blood) progenitors in the bone marrow and it function to differentiate hematopoietic stem cells, which are CD135 negative, from multipotent progenitors, which are CD135 positive. CD135 is a receptor tyrosine kinase typeIII for the cytokine Flt3 ligand and activat signaling through second messengers by binding to Flt3. Signaling through CD135 is important for lymphocyte development. The encoding gene CD135 is a proto-oncogene to which mutations happened will lead to cancer such as leukemia.Cancer ImmunotherapyImmune CheckpointImmunoTherapyTargeted Therapy

**Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins**

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E\_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481