

Anti-Complement C5a Antibody (8J841)

Product Details

Ig Type:	Rabbit IgG
Reactivity:	Human
Conjugation:	Unconjugated
Clone:	8J841
Purification:	Protein A

Applications

Application:	ELISA
Recommended	ELISA: 1:5000-1:10000

Properties

Stability & Storage:	Store at 2°C-8°C for 1 month. Store at -20°C or -80°C for 12 months. Avoid repeated freeze-thaw cycles. Preservative-Free.
Shipping:	Shipping with blue ice.

Antigen Details

Immunogen:	Recombinant Protein: Human Complement C5a Protein (TMPY-00653)
Antigen Species:	Human
Synonyms:	complement component 5

Research Background

C5a is a protein fragment released from complement component C5. This 74 amino acid peptide in humans is generated by the cleavage of C5a convertase on the C5 α -chain during the classical, alternative, and lectin pathways of complement activation. The structure of C5a includes a core region consisting of four, anti-parallel alpha-helices held together by three disulfide linkages and a structured C-terminal tail, and C5a is rapidly metabolised by carboxypeptidase B to a 73 amino acid low activity form, C5a des-Arg. C5a is an extremely potent proinflammatory mediator, as well as a potent chemotactic factor for neutrophils and other leukocytes. It causes histamine release, increases in vascular permeability, induces several cytokines production from leukocytes, enhances neutrophil-endothelial cell adhesion, and augments the humoral and cell-mediated immune response. C5a is quickly metabolised by carboxypeptidases, forming the less potent C5adesArg. Acting via a classical G protein-coupled receptor, CD88, C5a and C5adesArg exert a number of effects essential to the innate immune response, while their actions at the more recently discovered non-G protein-coupled receptor, C5L2 (or GPR77), remain unclear. The widespread expression of C5a receptors throughout the body allows C5a to elicit a broad range of effects. Thus, C5a has been found to be a significant pathogenic driver in a number of immuno-inflammatory diseases, making C5a inhibition an attractive therapeutic strategy. C5a is a strong chemoattractant and is involved in the recruitment of inflammatory cells such as neutrophils, eosinophils, monocytes, and T lymphocytes, in activation of phagocytic cells and release of granule-based enzymes and generation of oxidants, all of which may contribute to innate immune functions or tissue damage. Accordingly, the anaphylatoxin C5a is implicated in a variety of diseases such as rheumatoid arthritis, systemic lupus erythematosus, reperfusion injury, Alzheimer's disease, and sepsis.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481