

Anti-FGFR1 Antibody (7J437)

Product Details

Ig Type:	Mouse IgG2a
Reactivity:	Human
Conjugation:	Unconjugated
Clone:	7J437
Purification:	Protein A

Applications

Verified Activity:	1. Immunochemical staining of FGFR1 in pancreas with mouse monoclonal antibody (1:60, formalin-fixed paraffin embedded sections). 2. Immunochemical staining of FGFR1 in lung carcinoma with mouse monoclonal antibody (1:60, formalin-fixed paraffin embedded sections).
Application:	IHC-P
Recommended	IHC-P: 1:50-1:200

Properties

Stability & Storage:	Store at 2°C-8°C for 1 month. Store at -20°C or -80°C for 12 months. Avoid repeated freeze-thaw cycles. Preservative-Free.
Shipping:	Shipping with blue ice.

Antigen Details

Immunogen:	Recombinant Protein: Human FGFR1 / CD331 protein (TMPY-01011)
Antigen Species:	Human
Synonyms:	FGFR1 α (IIIC);fibroblast growth factor receptor 1;CD331;N-SAM;CEK;HBGFR;HRTFDS;FGFR-1;HH2;OGD;bFGF-R-1;BFGFR;KAL2;FLT2;FLT-2;FLG;FGFBR
Biology Area:	Cancer Drug Targets, Receptor Tyrosine Kinases (RTKs)

Research Background

FGFR1, also known as CD331, belongs to the fibroblast growth factor receptor subfamily where amino acid sequence is highly conserved between members and throughout evolution. FGFR family members differ from one another in their ligand affinities and tissue distribution. Fibroblast growth factors (FGFs) (FGF1 - 10 and 16 - 23) are mitogenic signaling molecules that have roles in angiogenesis, wound healing, cell migration, neural outgrowth and embryonic development. FGFs bind heparan sulfate glycosaminoglycans, which facilitates dimerization (activation) of FGF receptors. FGFR1 is a full-length representative protein consists of an extracellular region, composed of three immunoglobulin-like domains, a single hydrophobic membrane-spanning segment and a cytoplasmic tyrosine kinase domain. The extracellular portion of FGFR1 interacts with fibroblast growth factors, setting in motion a cascade of downstream signals, ultimately influencing mitogenesis and differentiation. This particular family member binds both acidic and basic fibroblast growth factors and is involved in limb induction. CD331 can be detected in astrocytoma, neuroblastoma and adrenal cortex cell lines. Some isoforms are detected in foreskin fibroblast cell lines, however isoform 17, isoform 18 and isoform 19 are not detected in these cells. Defects in FGFR1 are a cause of Pfeiffer syndrome, idiopathic hypogonadotropic hypogonadism, Kallmann syndrome type 2,

A DRUG SCREENING EXPERT

osteoglophonic dysplasia and trigonocephaly non-syndromic.Cancer ImmunotherapyImmune
CheckpointImmunotherapyTargeted Therapy

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only· Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481