

Anti-HGFR/c-Met Antibody-PE (6L502)

Product Details

Ig Type:	Rabbit IgG
Reactivity:	Human
Conjugation:	PE
Clone:	6L502
Purification:	Protein A

Applications

Verified Activity:	Profile of anti-human C-Met (HGF R) reactivity on HepG2 cells analyzed by flow cytometry. The Clone 6L502 and 4V580 monoclonal antibodies are not cross-blocking.
Application:	FCM
Recommended	5 µl/Test, 0.1 mg/ml

Properties

Stability & Storage:	Store at 2°C-8°C for 12 months, do not freeze. Keep away from direct sunlight.
Shipping:	Shipping with blue ice.

Antigen Details

Immunogen:	Recombinant Protein: Human HGFR / c-MET protein (TMPY-00718)
Antigen Species:	Human
Synonyms:	MET proto-oncogene, receptor tyrosine kinase
Biology Area:	Cancer Drug Targets, Receptor Tyrosine Kinases (RTKs)

Research Background

Hepatocyte growth factor receptor (HGFR), also known as c-Met or mesenchymal-epithelial transition factor (MET), is a receptor tyrosine kinase (RTK) that is overexpressed and/or mutated in a variety of malignancies. HGFR protein is produced as a single-chain precursor, and HGF is the only known ligand. Normal HGF/HGFR signaling is essential for embryonic development, tissue repair, or wound healing, whereas aberrantly active HGFR has been strongly implicated in tumorigenesis, particularly in the development of invasive and metastatic phenotypes. HGFR protein is a multifaceted regulator of growth, motility, and invasion, and is normally expressed by cells of epithelial origin. Preclinical studies suggest that targeting aberrant HGFR signaling could be an attractive therapy in cancer. Cancer Immunotherapy Immune Checkpoint Immunotherapy Targeted Therapy

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel: 781-999-4286 E_mail: info@targetmol.com Address: 34 Washington Street, Wellesley Hills, MA 02481