

Anti-CD38 Antibody (4R821)

Product Details

Ig Type:	Rabbit IgG
Reactivity:	Human
Conjugation:	Unconjugated
Clone:	4R821
Purification:	Protein A

Applications

Verified Activity:	1. Immunochemical staining of human CD38 in human tonsil with rabbit monoclonal antibody (1:200, formalin-fixed paraffin embedded sections).
	2. Immunochemical staining of human CD38 in human thymus with rabbit monoclonal antibody (1:200, formalin-fixed paraffin embedded sections).
	3. Immunochemical staining of human CD38 in human spleen with rabbit monoclonal antibody (1:200, formalin-fixed paraffin embedded sections).
Application:	IHC-P
Recommended	IHC-P: 1:100-1:500

Properties

Stability & Storage:	Store at 2°C-8°C for 1 month. Store at -20°C or -80°C for 12 months. Avoid repeated freeze-thaw cycles. Preservative-Free.
Shipping:	Shipping with blue ice.

Antigen Details

Immunogen:	Recombinant Protein: Human CD38 protein (TMPY-01023)
Antigen Species:	Human
Synonyms:	ADPRC1;ADPRC 1;T10;CD38 molecule
Biology Area:	Cancer Drug Targets

Research Background

The cluster of differentiation (CD) system is commonly used as cell markers in Immunophenotyping. Different kinds of cells in the immune system can be identified through the surface CD molecules associating with the immune function of the cell. There are more than 320 CD unique clusters and subclusters have been identified. Some of the CD molecules serve as receptors or ligands important to the cell through initiating a signal cascade which then alter the behavior of the cell. Some CD proteins do not take part in cell signal process but have other functions such as cell adhesion. Cluster of differentiation 38 (CD38), also known as ADP-ribosyl cyclase, is a glycoprotein found on the surface of many immune cells (white blood cells), including CD4+, CD8+, B and natural killer cells. It shares several characteristics with ADP-ribosyl cyclase 2 CD157. CD38 is a multifunctional ectoenzyme that catalyzes the synthesis and hydrolysis of cyclic ADP-ribose (cADPR) from NAD⁺ to ADP-ribose. It also functions in cell adhesion, signal transduction and calcium signaling. CD38 has been used as a prognostic marker in leukemia. It can also be used to identify plasma cells.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481