

Anti-CXCL1 Antibody (2U261)

Product Details

Ig Type:	Mouse IgG2a
Reactivity:	Human
Conjugation:	Unconjugated
Clone:	2U261
Purification:	Protein A

Applications

Application:	ELISA(Det)
--------------	------------

Properties

Stability & Storage:	Store at 2°C-8°C for 1 month. Store at -20°C or -80°C for 12 months. Avoid repeated freeze-thaw cycles. Preservative-Free.
Shipping:	Shipping with blue ice.

Antigen Details

Immunogen:	Recombinant Protein: Human CXCL1 Protein (TMPY-02868)
Antigen Species:	Human
Synonyms:	GROa;NAP-3;CYP21;GRO1;CA21H;MGSA-a;CAH1;MGSA;SCYB1;P450c21B;CYP21B;chemokine (C-X-C motif) ligand 1 (melanoma growth stimulating activity, α);chemokine (C-X-C motif) ligand 1 (melanoma growth stimulating activity, alpha);CPS1

Research Background

The Chemokine (C-X-C motif) Ligand 1, CXCL1, is a small cytokine belonging to the CXC chemokine family that was previously called GRO1 oncogene, GRO γ , KC, Neutrophil-activating protein 3 (NAP-3) and melanoma growth stimulating activity, alpha (MSGA-a). CXCL1 already known to be important in osteoarthritis (OA), as a novel target gene of transcription factor AP-2 γ in chondrocytes and support the important role of AP-2 γ in cartilage. CXCL1 is a potent neutrophil chemoattractant with recognized roles in angiogenesis and inflammation. CXCL1 is a novel immediate PTH/PTHrP-responsive gene. CXCL1 may act as a chemoattractant for osteoclast precursors. CXCL1 may also have important pro-nociceptive effects via its direct actions on sensory neurons, and may induce long-term changes that involve protein synthesis. CXCL1 plays a critical nonredundant role in the development of experimental Lyme arthritis and carditis via CXCR2-mediated recruitment of neutrophils into the site of infection. CXCL1 functions through CXCR2 to transactivate the EGFR by proteolytic cleavage of HB-EGF, leading to activation of MAPK signalling and increased proliferation of epithelial ovarian cancer (EOC) cells. It might limit tumor growth by reinforcing senescence early in tumorigenesis. Thus, CXCL1 plays a role in spinal cord development by inhibiting the migration of oligodendrocyte precursors and is involved in the processes of angiogenesis, inflammation, wound healing, and tumorigenesis.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481