

Anti-CD73 Antibody (8K319)

Product Details

Ig Type:	Rabbit IgG
Reactivity:	Human
Conjugation:	Unconjugated
Clone:	8K319
Purification:	Protein A

Applications

Verified Activity:	Immunofluorescence staining of NT5E-3 in MCF7 cells. Cells were fixed with 4% PFA, blocked with 10% serum, and incubated with rabbit anti-human NT5E-3 monoclonal antibody (dilution ratio 1:60) at 4°C overnight. Then cells were stained with the Alexa Fluor®488-conjugated Goat Anti-rabbit IgG secondary antibody (green). Positive staining was localized to Cytoplasm .
Application:	ELISA, ICC/IF
Recommended	ELISA: 1:5000-1:10000; ICC-IF: 1:20-1:100

Properties

Stability & Storage:	Store at 2°C-8°C for 1 month. Store at -20°C or -80°C for 12 months. Avoid repeated freeze-thaw cycles. Preservative-Free.
Shipping:	Shipping with blue ice.

Antigen Details

Immunogen:	Recombinant Protein: Human CD73 / NT5E protein (TMPY-01851)
Antigen Species:	Human
Synonyms:	5'-nucleotidase, ecto (CD73);NTE;eNT;NT;CD73;E5NT;CALJA;NT5;eN
Biology Area:	Phosphatases and Regulators

Research Background

5'-nucleotidase, also known as NT5E, NTE, and CD73, is a cell membrane protein that belongs to the 5'-nucleotidase family. CD73 is a glycosylphosphatidylinositol (GPI) anchored purine salvage enzyme expressed on the surface of human T and B lymphocytes. CD73 catalyzes the conversion of purine and pyrimidine ribo- and deoxyribonucleoside monophosphates to the corresponding nucleosides. CD73 serves as a costimulatory molecule in activating T cells. CD73 generated adenosine functions in cell signaling in many physiologic systems, including intestinal epithelium, ischemic myocardium, and cholinergic synapses. CD73 might mediate lymphocyte-stromal cell interactions or condition the local microenvironment to facilitate lymphocyte development and/or function. In CD73-depleted cells, surface levels of the leukocyte adhesion molecules ICAM-1, VCAM-1, and E-selectin increase. CD73 produces extracellular adenosine, which then acts on G protein-coupled purinergic receptors to induce cellular responses. CD73 has also been reported to regulate the expression of pro-inflammatory molecules in mouse endothelium.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only. Not for Human or Veterinary or Therapeutic Use

Tel: 781-999-4286 E_mail: info@targetmol.com Address: 34 Washington Street, Wellesley Hills, MA 02481