

Anti-CD20 Antibody (9C260)

Product Details

Ig Type:	Mouse IgG1
Reactivity:	Human, Cynomolgus
Conjugation:	Unconjugated
Clone:	9C260
Purification:	Protein A

Applications

Verified Activity:	1. Immunochemical staining CD20(L) in human tonsil with mouse monoclonal antibody (1:30, formalin-fixed paraffin embedded sections).
	2. Immunochemical staining CD20(L) in human spleen with mouse monoclonal antibody (1:30, formalin-fixed paraffin embedded sections).
	3. Immunochemical staining CD20 in cynomolgus spleen with mouse monoclonal antibody at 1:30 dilution, formalin-fixed paraffin embedded sections.
	4. Immunochemical staining CD20 in cynomolgus lymph node with mouse monoclonal antibody at 1:30 dilution, formalin-fixed paraffin embedded sections.
Application:	IHC-P
Recommended	IHC-P: 1:20-1:80

Properties

Stability & Storage:	Store at 2°C-8°C for 1 month. Store at -20°C or -80°C for 12 months. Avoid repeated freeze-thaw cycles. Preservative-Free.
Shipping:	Shipping with blue ice.

Antigen Details

Immunogen:	Recombinant Protein: Human CD20 / MS4A1 Protein (TMPY-02819)
Antigen Species:	Human
Synonyms:	membrane-spanning 4-domains, subfamily A, member 1;CVID5;S7;CD20;MS4A2;MS4A1;B1;LEU-16;Bp35
Biology Area:	Cancer Drug Targets

Research Background

CD20 (membrane-spanning 4-domains, subfamily A, member 1), also known as MS4A1, is a member of the membrane-spanning 4A gene family. Members of this nascent protein family are characterized by common structural features and similar intron/exon splice boundaries and display unique expression patterns among hematopoietic cells and nonlymphoid tissues. CD20 / MS4A1 is expressed in all stages of B cell development except the first and last. CD20 / MS4A1 is present from pre-pre B cells through memory cells, but not on either pro-B cells or plasma cells. It is a B-lymphocyte surface molecule that plays a role in the development and differentiation of B-cells into plasma cells. CD20 / MS4A1 may be involved in the regulation of B-cell activation and proliferation. Defects in CD20 / MS4A1 are the cause of immunodeficiency common variable type 5 (CVID5). CVID5 is a primary immunodeficiency characterized by antibody deficiency, hypogammaglobulinemia, recurrent bacterial infections,

A DRUG SCREENING EXPERT

and an inability to mount an antibody response to antigen. The defect results from a failure of B-cell differentiation and impaired secretion of immunoglobulins; the numbers of circulating B-cells are usually in the normal range but can be very low. Cancer Immunotherapy Immune Checkpoint Immunotherapy Targeted Therapy

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel: 781-999-4286 E_mail: info@targetmol.com Address: 34 Washington Street, Wellesley Hills, MA 02481