

## Anti-CEACAM5 Antibody (2J586)

### Product Details

Ig Type:	Rabbit IgG
Reactivity:	Human
Conjugation:	Unconjugated
Clone:	2J586
Purification:	Protein A

### Applications

Verified Activity:	1. Immunochemical staining of human CEACAM5 in human rectal cancer with rabbit monoclonal antibody (1:200, formalin-fixed paraffin embedded sections). Positive staining was localized to intestinal gland.
	2. Immunochemical staining of human CEACAM5 in human colon carcinoma with rabbit monoclonal antibody (1:200, formalin-fixed paraffin embedded sections). Positive staining was localized to intestinal gland.
Application:	IHC-P
Recommended	IHC-P: 1:100-1:500

### Properties

Stability & Storage:	Store at 2°C-8°C for 1 month. Store at -20°C or -80°C for 12 months. Avoid repeated freeze-thaw cycles. Preservative-Free.
Shipping:	Shipping with blue ice.

### Antigen Details

Immunogen:	Recombinant Protein: Human CEACAM5 / CD66e protein (TMPY-01647)
Antigen Species:	Human
Synonyms:	carcinoembryonic antigen-related cell adhesion molecule 5;CD66e;CEA
Biology Area:	Cancer Drug Targets

### Research Background

CEACAM5, also known as CEA or CD66e, belongs to the large CEACAM subfamily of the immunoglobulin superfamily. CEACAM5 is expressed primarily by epithelial cells, and is synthesized as a glycoprotein with an MW of 180 kDa comprising 60% carbohydrate. CEACAM5 contains one Ig-like V-type domain at the N-terminus, followed by six Ig-like C2-type domains and a GPI anchor, and exists as a homodimer. CEACAM5 and CEACAM6 are overexpressed in many cancers and are associated with adhesion and invasion. CEACAM5 can mediate cell-cell adhesion through homotypic and heterotypic interactions. It functions as a homotypic intercellular adhesion molecule and serves as a widely used tumor marker, since it is expressed at higher levels in tumorous tissues than in corresponding normal tissues. CEACAM5 has also been shown to contribute to tumorigenicity by inhibiting cellular differentiation. In addition, CEACAM5 is identified as the host receptor for the Dr family of adhesins of E.Coli, and the binding of E.coli Dr adhesins leads to dissociation of the CEACAM5 homodimer. Cancer Immunotherapy Immune Checkpoint Immunotherapy Targeted Therapy

**Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins**

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E\_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481