

## Anti-S100A9 Antibody (2K33)

### Product Details

Ig Type:	Rabbit IgG
Reactivity:	Human
Conjugation:	Unconjugated
Clone:	2K33
Purification:	Protein A

### Applications

	Anti-S100A9 rabbit monoclonal antibody at 1:500 dilution. -Lane A: HL-60 Whole Cell lysate. -Lysates/proteins at 30 µg per lane. -Secondary
Verified Activity:	-Goat Anti-Rabbit IgG H&L (Dylight800) at 1/10000 dilution. -Developed using the Odyssey technique. -Performed under reducing conditions. -Predicted band size:13 kDa. -Observed band size:15 kDa
Application:	ELISA,WB
Recommended	WB: 1:500-1:2000; ELISA: 1:5000-1:10000

### Properties

Stability & Storage:	Store at 2°C-8°C for 1 month. Store at -20°C or -80°C for 12 months. Avoid repeated freeze-thaw cycles. Preservative-Free.
Shipping:	Shipping with blue ice.

### Antigen Details

Immunogen:	Recombinant Protein: Human S100A9 Protein (TMPY-01330)
Antigen Species:	Human
Synonyms:	p14;GAGB;BEE22;MRP14;Cagb;S100 calcium binding protein A9;L1Ag;AW546964;60B8Ag
Biology Area:	Calcium-binding Proteins and Related Molecules

### Research Background

S100 protein is a family of low molecular weight protein found in vertebrates characterized by two EF-hand calcium-binding motifs. There are at least 21 different S100 proteins, and the name is derived from the fact that the protein is 100% soluble in ammonium sulfate at neutral pH. Most S100 proteins are disulfide-linked homodimer, and is normally present in cells derived from the neural crest, chondrocytes, macrophages, dendritic cells, etc. S100 proteins have been implicated in a variety of intracellular and extracellular functions. They are involved in regulation of protein phosphorylation, transcription factors, the dynamics of cytoskeleton constituents, enzyme activities, cell growth and differentiation, and the inflammatory response. Protein S100-A9, also known as S100 calcium binding protein A9, S100A9, and CAGB, is a member of the S-100 family. S100A9 is expressed by macrophages in acutely inflamed tissues and in chronic inflammation. It is also expressed in epithelial cells constitutively or induced during

## A DRUG SCREENING EXPERT

---

dermatoses. S100A9 is a calcium-binding protein. It has anti-microbial activity towards bacteria and fungi. The anti-microbial and proapoptotic activity of S100A9 is inhibited by zinc ions. S100A9 plays a role in the development of endotoxic shock in response to bacterial lipopolysaccharide (LPS). It promotes tubulin polymerization when unphosphorylated. It also promotes phagocyte migration and infiltration of granulocytes at sites of wounding. S100A9 plays a role as a proinflammatory mediator in acute and chronic inflammation and up-regulates the release of IL8 and cell-surface expression of ICAM1.

**Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins**

This product is for Research Use Only· Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E\_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481