

Anti-S100A9 Antibody (6Z270)

Product Details

| | |
|---------------|--------------|
| Ig Type: | Rabbit IgG |
| Reactivity: | Human |
| Conjugation: | Unconjugated |
| Clone: | 6Z270 |
| Purification: | Protein A |

Applications

| | |
|--------------------|---|
| Verified Activity: | Immunochemical staining of human S100A9 in human spleen with rabbit monoclonal antibody (1:200, formalin-fixed paraffin embedded sections). |
| Application: | IHC-P |
| Recommended | IHC-P: 1:100-1:500 |

Properties

| | |
|----------------------|--|
| Stability & Storage: | Store at 2°C-8°C for 1 month. Store at -20°C or -80°C for 12 months. Avoid repeated freeze-thaw cycles. Preservative-Free. |
| Shipping: | Shipping with blue ice. |

Antigen Details

| | |
|------------------|--|
| Immunogen: | Recombinant Protein: Human S100A9 protein (TMPY-01357) |
| Antigen Species: | Human |
| Synonyms: | BEE22;L1Ag;MRP14;Cagb;GAGB;S100 calcium binding protein A9;AW546964;p14;60B8Ag |
| Biology Area: | Calcium-binding Proteins and Related Molecules |

Research Background

S100 protein is a family of low molecular weight protein found in vertebrates characterized by two EF-hand calcium-binding motifs. There are at least 21 different S100 proteins, and the name is derived from the fact that the protein is 100% soluble in ammonium sulfate at neutral pH. Most S100 proteins are disulfide-linked homodimer, and is normally present in cells derived from the neural crest, chondrocytes, macrophages, dendritic cells, etc. S100 proteins have been implicated in a variety of intracellular and extracellular functions. They are involved in regulation of protein phosphorylation, transcription factors, the dynamics of cytoskeleton constituents, enzyme activities, cell growth and differentiation, and the inflammatory response. Protein S100-A9, also known as S100 calcium binding protein A9, S100A9, and CAGB, is a member of the S-100 family. S100A9 is expressed by macrophages in acutely inflamed tissues and in chronic inflammation. It is also expressed in epithelial cells constitutively or induced during dermatoses. S100A9 is a calcium-binding protein. It has anti-microbial activity towards bacteria and fungi. The anti-microbial and proapoptotic activity of S100A9 is inhibited by zinc ions. S100A9 plays a role in the development of endotoxic shock in response to bacterial lipopolysaccharide (LPS). It promotes tubulin polymerization when unphosphorylated. It also promotes phagocyte migration and infiltration of granulocytes at sites of wounding. S100A9 plays a role as a proinflammatory mediator in acute and chronic inflammation and up-regulates the release of IL8 and cell-surface expression of ICAM1.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481