

Anti-CD200RLa Antibody (8K476)

Product Details

| | |
|---------------|--------------|
| Ig Type: | Rabbit IgG |
| Reactivity: | Human |
| Conjugation: | Unconjugated |
| Clone: | 8K476 |
| Purification: | Protein A |

Applications

| | |
|--------------|-----------------------|
| Application: | ELISA |
| Recommended | ELISA: 1:5000-1:10000 |

Properties

| | |
|----------------------|--|
| Stability & Storage: | Store at 2°C-8°C for 1 month. Store at -20°C or -80°C for 12 months. Avoid repeated freeze-thaw cycles. Preservative-Free. |
| Shipping: | Shipping with blue ice. |

Antigen Details

| | |
|------------------|--|
| Immunogen: | Recombinant Protein: Human CD200R1L protein (TMPY-01591) |
| Antigen Species: | Human |
| Synonyms: | CD200 receptor 1 like;CD200RLa |

Research Background

Cell surface glycoprotein CD2 receptor 2, also known as Cell surface glycoprotein CD2 receptor 1-like, Cell surface glycoprotein OX2 receptor 2, CD2 receptor-like 2, CD2R1a, CD2R1L and CD2R2, is a single-pass type I membrane protein which belongs to the CD2R family. CD2R1L / CD2R2. It contains one Ig-like C2-type (immunoglobulin-like) domain and one Ig-like V-type (immunoglobulin-like) domain. CD2 is a transmembrane protein delivering immunoregulatory signals after engagement of CD2R. A family of CD2Rs exist (CD2R1, CD2R2, CD2R3, CD2R4) with different tissue expression and functional activity. In the presence of anti-CD2R2 / CD2R3 monoclonal antibodies (mAbs), bone-marrow cells cultured in the presence of (interleukin [IL]-4+granulocyte-macrophage colony-stimulating factor) differentiate into dendritic cells (DCs), which induce CD4+CD25+ Treg. Interaction between the relatively ubiquitously expressed molecule CD2 and one of its receptors, CD2R1, resulted in direct suppression of alloreactivity, engagement of alternate receptors led instead to altered differentiation of dendritic cells (DCs) from marrow precursors, which could in turn foster development of Foxp3(+) regulatory T cells. Unlike anti-CD2R1, anti-CD2R2 both promotes development of DCs with capacity to induce Treg and directly augments thymocyte production of Treg.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only. Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481