

Anti-IL-2 Antibody (9Z926)

Product Details

Ig Type:	Rabbit IgG
Reactivity:	Human
Conjugation:	Unconjugated
Clone:	9Z926
Purification:	Protein A

Applications

Application:	ELISA
Recommended	ELISA: 1:5000-1:10000

Properties

Stability & Storage:	Store at 2°C-8°C for 1 month. Store at -20°C or -80°C for 12 months. Avoid repeated freeze-thaw cycles. Preservative-Free.
Shipping:	Shipping with blue ice.

Antigen Details

Immunogen:	Recombinant Protein: Human IL2 Protein
Antigen Species:	Human
Synonyms:	IL-2;interleukin 2;Interleukin-2;TCGF;lymphokine
Biology Area:	Neuroinflammation

Research Background

Interleukin-2, also known as a T-cell growth factor, TCGF, Aldesleukin, and IL2, is a secreted protein that belongs to the IL-2 family. Interleukin-2 / IL-2 was the first interleukin molecule to be discovered. Interleukin-2 / IL-2 molecule was first purified to homogeneity by immunoaffinity chromatography by Kendall Smith and his team at Dartmouth Medical School. Interleukin-2 / IL-2 was also the first cytokine shown to mediate its effects via a specific IL-2 receptor, and it was also the first interleukin to be cloned and expressed from a complementary DNA (cDNA) library. Interleukin-2 / IL-2 was designated number 2 because Smith's data at the time indicated that IL-1, produced by macrophages, facilitates IL-2 production by T lymphocytes (T cells). Interleukin-2 / IL-2 is produced by T-cells in response to antigenic or mitogenic stimulation, this protein is required for T-cell proliferation and other activities crucial to regulation of the immune response. Interleukin-2 / IL-2 is normally produced by the body during an immune response. When environmental substances (molecules or microbes) gain access to the body, these substances (termed antigens) are recognized as foreign by antigen receptors that are expressed on the surface of lymphocytes. Antigen binding to the T cell receptor (TCR) stimulates the secretion of Interleukin-2 / IL-2 and the expression of IL-2 receptors IL-2R. The IL-2 / IL-2R interaction then stimulates the growth, differentiation, and survival of antigen-selected cytotoxic T cells via the activation of the expression of specific genes. Interleukin-2 / IL-2 can stimulate B-cells, monocytes, lymphokine-activated killer cells, natural killer cells, and glioma cells. The World Reference Standard for Interleukin-2 / IL-2 is produced by the National Institute of Biological Standards and Control in the UK. A recombinant form of Interleukin-2 / IL-2 for clinical use is manufactured by Chiron Corporation with the brand name Proleukin. It has been approved by the Food and Drug Administration (FDA) for the treatment of cancers

A DRUG SCREENING EXPERT

(malignant melanoma, renal cell cancer), and is in clinical trials for the treatment of chronic viral infections, and as a booster (adjuvant) for vaccines. The use of Interleukin-2 / IL-2 in HIV therapy is ineffective. Cancer Immunotherapy Immune Checkpoint Immunotherapy Targeted Therapy

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel: 781-999-4286 E_mail: info@targetmol.com Address: 34 Washington Street, Wellesley Hills, MA 02481