

## Anti-CD19 Antibody-FITC (1T602)

### Product Details

Ig Type:	Mouse IgG1
Reactivity:	Human
Conjugation:	FITC
Clone:	1T602
Purification:	Protein A

### Applications

Verified Activity:	Flow cytometric analysis of Human CD19 expression on human whole blood lymphocytes. Cells were stained with FITC-conjugated anti-Human CD19 and PE-conjugated anti-Human CD3. The fluorescence histograms were derived from gated events with the forward and side light-scatter characteristics of viable lymphocytes.
Application:	FCM
Recommended	10 µl/Test, 0.1 mg/ml

### Properties

Stability & Storage:	Store at 2°C-8°C for 12 months, do not freeze. Keep away from direct sunlight.
Shipping:	Shipping with blue ice.

### Antigen Details

Immunogen:	Recombinant Protein: Human CD19 Protein (TMPY-01949)
Antigen Species:	Human
Synonyms:	B4;CD19 molecule;CVID3
Biology Area:	Cancer Drug Targets

### Research Background

The cluster of differentiation (CD) system is commonly used as cell markers in Immunophenotyping. Different kinds of cells in the immune system can be identified through the surface CD molecules associating with the immune function of the cell. There are more than 320 CD unique clusters and subclusters have been identified. Some of the CD molecules serve as receptors or ligands important to the cell through initiating a signal cascade which then alter the behavior of the cell. Some CD proteins do not take part in cell signal process but have other functions such as cell adhesion. Cluster of differentiation 19 (CD19) is a member of CD system. CD19 is a cell surface molecule that assembles with the antigen receptor of B-cells. This results in a descent in the threshold for antigen receptor-dependent stimulation. A simplified view holds that the ability of B-cells to respond to the various antigens in a specific and sensitive manner is achieved in the presence of low-affinity antigen receptors. CD19 primarily acts as a B-cell co-receptor in conjunction with CD21 and CD81. The formation of the receptor complex is induced by antigen and CD19, induced by exogenous antigen, has been found cytoplasmic tail phosphorylated and bind to slg. Cancer Immunotherapy/Immune Checkpoint/Immunotherapy/Targeted Therapy

**Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins**

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E\_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481