

## Anti-CD44 Antibody (1S404)

## Product Details

Ig Type:	Rabbit IgG
Reactivity:	Human
Conjugation:	Unconjugated
Clone:	1S404
Purification:	Protein A

## Applications

Verified Activity:	Anti-CD44 rabbit monoclonal antibody at 1:500 dilution. -Lane A: HL60 Whole Cell Lysate. -Lane B: Hela Whole Cell lysate. -Lysates/proteins at 30 µg per lane. -Secondary -Goat Anti-Mouse IgG (H+L)/HRP at 1/10000 dilution. -Developed using the ECL technique. -Performed under reducing conditions. -Predicted band size:82 kDa. -Observed band size:82 kDa(We are unsure as to the identity of these extra bands.)
Application:	ELISA,WB
Recommended	WB: 1:500-1:2000; ELISA: 1:5000-1:10000

## Properties

Stability & Storage:	Store at 2°C-8°C for 1 month. Store at -20°C or -80°C for 12 months. Avoid repeated freeze-thaw cycles. Preservative-Free.
Shipping:	Shipping with blue ice.

## Antigen Details

Immunogen:	Recombinant Protein: Human CD44 protein (TMPY-02033)
Antigen Species:	Human
Synonyms:	HUTCH-I;ECMR-III;MDU3;MDU2;CSPG8;MIC4;HCELL;CDW44;CD44 molecule (Indian blood group);Pgp1;IN;MC56;LHR

## Research Background

CD44 is a type I transmembrane protein and a member of the cartilage link protein family. It is involved in cell-cell and cell-matrix interactions and signal transduction. Several CD44 ligands have been identified. The most extensively characterized ligand for CD44 is hyaluronan, a component of the extracellular matrix. CD44 protein is expressed on the majority of immune cells. The binding of CD44 to hyaluronan is induced on T lymphocytes after activation by antigen and on monocytes after stimulation by inflammatory agents. Under inflammatory conditions, CD44 on endothelial cells presents hyaluronan to CD44 on activated T lymphocytes and mediates a rolling interaction under flow conditions. Perturbations of the hyaluronan-CD44 interaction at the plasma membrane by various antagonists result in attenuation of receptor tyrosine kinase and transporter activities and inhibition of tumor progression in vivo. CD44 is known to interact with the ezrin family (ERM family) members and form a complex

## A DRUG SCREENING EXPERT

---

plays diverse roles within both normal and abnormal cells, particularly cancer cells. CD44 and ezrin and their respective complex have properties suggesting that they may be important in the process of tumour-endothelium interactions, cell migrations, cell adhesion, tumour progression and metastasis.

**Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins**

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E\_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481