

Anti-CD164 Antibody (2K50)

Product Details

Ig Type:	Mouse IgG1
Reactivity:	Human
Conjugation:	Unconjugated
Clone:	2K50
Purification:	Protein A

Applications

Verified Activity:	1. Immunochemical staining of human CD164 in human prostatic carcinoma with mouse monoclonal antibody (1:60, formalin-fixed paraffin embedded sections).
	2. Immunochemical staining of human CD164 in human pancreatic carcinoma with mouse monoclonal antibody (1:60, formalin-fixed paraffin embedded sections).
	3. Flow cytometric analysis of Human CD164 expression on human whole blood monocytes. Cells were stained with purified anti-Human CD164, then a FITC-conjugated second step antibody. The fluorescence histograms were derived from gated events with the forward and side light-scatter characteristics of viable monocytes.
Application:	ELISA,FCM,IHC-P
Recommended	ELISA: 1:1000-1:2000; IHC-P: 1:50-1:200; FCM: 1:25-1:100

Properties

Stability & Storage:	Store at 2°C-8°C for 1 month. Store at -20°C or -80°C for 12 months. Avoid repeated freeze-thaw cycles. Preservative-Free.
Shipping:	Shipping with blue ice.

Antigen Details

Immunogen:	Recombinant Protein: Human CD164 Protein (TMPY-02067)
Antigen Species:	Human
Synonyms:	CD164 molecule, sialomucin

Research Background

Sialomucin core protein 24 also known as endolyn or CD164 (cluster of differentiation 164) is a novel 80- to 90-kD mucin-like molecule expressed by human CD34+ hematopoietic progenitor cells. Isoform 1 and isoform 3 of CD164 are expressed in hematopoietic and non-hematopoietic tissues. Isoform 1 is expressed by prostate cancer tumors and prostate cancer cell lines. The expression is greater in bone metastases than in primary tumors. Expression in osseous metastasis is greater than that in soft tissue metastasis. Isoform 2 of CD164 is expressed in the small intestine, colon, lung, thyroid and colorectal and pancreatic adenocarcinoma. Isoform 4 is expressed by both hematopoietic progenitor cells and bone marrow stromal cells. CD164 belongs to the CD164 family. The cluster of differentiation (cluster of designation) (often abbreviated as CD) is a protocol used for the identification and investigation of cell surface molecules present on white blood cells initially but found in almost any kind of cell of the body, providing targets for immunophenotyping of cells. CD164 may play an important role in prostate cancer metastasis and the infiltration of bone marrow by cancer cells. CD164 promotes myogenesis by enhancing CXCR4-

A DRUG SCREENING EXPERT

dependent cell motility. This protein positively regulates myoblast migration and promotes myoblast fusion into myotubes. CD164 may play a key role in hematopoiesis by facilitating the adhesion of CD34+ cells to bone marrow stroma and by negatively regulating CD34+hematopoietic progenitor cell growth.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481