

Anti-CD164 Antibody-FITC (9R274)

Product Details

Ig Type:	Rabbit IgG
Reactivity:	Human
Conjugation:	FITC
Clone:	9R274
Purification:	Protein A

Applications

Verified Activity:	Flow cytometric analysis of Human CD164 expression on human whole blood monocytes. Cells were stained with FITC-conjugated anti-Human CD164. The fluorescence histograms were derived from gated events with the forward and side light-scatter characteristics of viable monocytes.
Application:	FCM
Recommended	10 µl/Test, 0.1 mg/ml

Properties

Stability & Storage:	Store at 2°C-8°C for 12 months, do not freeze. Keep away from direct sunlight. Sodium azide is toxic to cells and should be disposed of properly. Flush with large volumes of water during disposal.
Shipping:	Shipping with blue ice.

Antigen Details

Immunogen:	Recombinant Protein: Human CD164 Protein (TMPY-02067)
Antigen Species:	Human
Synonyms:	CD164 molecule, sialomucin

Research Background

Sialomucin core protein 24 also known as endolyn or CD164 (cluster of differentiation 164) is a novel 80- to 90-kD mucin-like molecule expressed by human CD34+ hematopoietic progenitor cells. Isoform 1 and isoform 3 of CD164 are expressed in hematopoietic and non-hematopoietic tissues. Isoform 1 is expressed by prostate cancer tumors and prostate cancer cell lines. The expression is greater in bone metastases than in primary tumors. Expression in osseous metastasis is greater than that in soft tissue metastasis. Isoform 2 of CD164 is expressed in the small intestine, colon, lung, thyroid and colorectal and pancreatic adenocarcinoma. Isoform 4 is expressed by both hematopoietic progenitor cells and bone marrow stromal cells. CD164 belongs to the CD164 family. The cluster of differentiation (cluster of designation) (often abbreviated as CD) is a protocol used for the identification and investigation of cell surface molecules present on white blood cells initially but found in almost any kind of cell of the body, providing targets for immunophenotyping of cells. CD164 may play an important role in prostate cancer metastasis and the infiltration of bone marrow by cancer cells. CD164 promotes myogenesis by enhancing CXCR4-dependent cell motility. This protein positively regulates myoblast migration and promotes myoblast fusion into myotubes. CD164 may play a key role in hematopoiesis by facilitating the adhesion of CD34+ cells to bone marrow stroma and by negatively regulating CD34+hematopoietic progenitor cell growth.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481