

Anti-IL-22 Antibody (6H655)

Product Details

Ig Type:	Mouse IgG3
Reactivity:	Human
Conjugation:	Unconjugated
Clone:	6H655
Purification:	Protein A

Applications

Application:	ELISA
Recommended	ELISA: 1:1000-1:2000

Properties

Stability & Storage:	Store at 2°C-8°C for 1 month. Store at -20°C or -80°C for 12 months. Avoid repeated freeze-thaw cycles. Preservative-Free.
Shipping:	Shipping with blue ice.

Antigen Details

Immunogen:	Recombinant Protein: Human IL22 / Interleukin 22 Protein (TMPY-03490)
Antigen Species:	Human
Synonyms:	TIFIL-23;interleukin 22;IL-D110;IL-TIF;ILTIF;IL-22;zcyto18;TIFa

Research Background

IL22 is a member of a group of cytokines called the IL-1 family of IL-1 superfamily (including IL-19, IL-2, IL-24, and IL-26), a class of potent mediators of cellular inflammatory responses. It shares the use of IL-1R2 in cell signaling with other members of this family, IL-1, IL-26, IL-28A/B, and IL-29. IL22 is produced by activated DC and T cells and initiates innate immune responses against bacterial pathogens especially in epithelial cells such as respiratory and gut epithelial cells. IL22 along with IL-17 is rapidly produced by splenic LTI-like cells and can be also produced by Th17 cells and likely plays a role in the coordinated response of both adaptive and innate immune systems. IL22 biological activity is initiated by binding to a cell-surface complex composed of IL-22R1 and IL-1R2 receptor chains and further regulated by interactions with a soluble binding protein, IL-22BP, which shares sequence similarity with an extracellular region of IL-22R1 (SIL-22R1). IL22 and IL-1 receptor chains play a role in cellular targeting and signal transduction to selectively initiate and regulate immune responses. IL22 can contribute to immune disease through the stimulation of inflammatory responses, S1s, and defensins. IL22 also promotes hepatocyte survival in the liver and epithelial cells in the lung and gut similar to IL-1. In some contexts, the pro-inflammatory versus tissue-protective functions of IL22 are regulated by the often co-expressed cytokine IL-17A.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481