

Anti-BST2 Antibody (2B626)

Product Details

Ig Type:	Mouse IgG1
Reactivity:	Human
Conjugation:	Unconjugated
Clone:	2B626
Purification:	Protein A

Applications

Verified Activity:	Immunochemical staining of human BST2 in human liver with mouse monoclonal antibody (1:60, formalin-fixed paraffin embedded sections).
Application:	IHC-P
Recommended	IHC-P: 1:50-1:200

Properties

Stability & Storage:	Store at 2°C-8°C for 1 month. Store at -20°C or -80°C for 12 months. Avoid repeated freeze-thaw cycles. Preservative-Free.
Shipping:	Shipping with blue ice.

Antigen Details

Immunogen:	Recombinant Protein: Human BST2 Protein (TMPY-03235)
Antigen Species:	Human
Synonyms:	DAMP-1;Bst-2;CD317;2310015I10Rik;GREG;C87040;bone marrow stromal cell antigen 2

Research Background

BST2 was frequently overexpressed in GC tissues compared with the adjacent non-tumorous tissues, and high BST2 expression was correlated with tumor stage and lymphatic metastasis. Furthermore, in vitro experiments demonstrated that knockdown of BST2 by siRNA inhibited cell proliferation, induced apoptosis and repressed cell motility in GC cells. In addition, the pro-tumor function of BST2 in GC was mediated partly through the NF- κ B signaling. BST2 possesses the oncogenic potential in GC by regulating the proliferation, apoptosis, and migratory ability of GC cells, thereby BST2 could be a potential therapeutic target for the treatment of GC. IFN (interferon)-induced BST2 recruits the E3 ubiquitin ligase MARCH8 to catalyze the K27-linked ubiquitination of MAVS for CALCOCO2-directed autophagic degradation, hence inhibiting DDX58-mediated type I interferon signaling through a negative feedback loop. BST2 is a host protein with dual functions in response to viral infections: it traps newly assembled enveloped virions at the plasma membrane in infected cells, and it induces NF- κ B activity, especially in the context of retroviral assembly. BST2 may induce or amplify proinflammatory signaling during Ebola virus infection, potentially contributing to the dysregulated cytokine response that is a hallmark of Ebola virus disease.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481