

## Anti-SARS-CoV/SARS-CoV-2 Nucleocapsid Antibody (9Z26)

### Product Details

Ig Type:	Mouse IgG1
Conjugation:	Unconjugated
Clone:	9Z26
Purification:	Protein A

### Applications

Verified Activity:	1. Anti-SARS-NP mouse monoclonal antibody at 1:1000 dilution. -Lane A: SARS-CoV NP Protein (30ng) -Lane B: SARS-CoV NP Protein (5ng) -Lane C: SARS-CoV-2 (2019-nCoV) NP Protein (30ng) -Lane D: SARS-CoV-2 (2019-nCoV) NP Protein (5ng) -Secondary -Goat Anti-Mouse IgG (H+L)/HRP at 1/10000 dilution. -Developed using the ECL technique. -Performed under reducing conditions.
	2. Immunochemical analysis of Nucleocapsid overexpressed Hek293 Cells were stained with purified anti-SARS-CoV-2 Nucleocapsid Mouse Mab, then a HRP-conjugated second step antibody. (Validation Experiment)
	3. Anti-SARS-CoV / SARS-CoV-2 Nucleocapsid mouse monoclonal antibody at 1:5000 dilution. -Lane A: SARS-COV-2 Nucleocapsid (YP_009724397.2, WT) overexpressed HEK293 Whole Cell Lysate. -Lane B: HEK293 Whole Cell lysate. -Lysates/proteins at 10 µg per lane. -Secondary -Goat Anti-Mouse IgG (H+L)/HRP at 1/10000 dilution. -Developed using the ECL technique. -Performed under reducing condition(Validation Experiment)
	4. Flow cytometric analysis of SARS-COV-2 Nucleocapsid overexpressed HEK293 Cells were stained with purified anti-SARS-COV-2 Nucleocapsid Mouse MAb, then a FITC-conjugated second step antibody. The fluorescence histograms were derived from gated events with the forward and side light-scatter characteristics of intact cells. (Validation Experiment)
Application:	ELISA,FCM,IHC-P,WB
Recommended	WB: 1:1000-1:10000; ELISA: 1:1000-1:2000; IHC-P: 1:1000-1:5000; FCM: 1:25-1:100

### Properties

Stability & Storage:	Store at 2°C-8°C for 1 month. Store at -20°C or -80°C for 12 months. Avoid repeated freeze-thaw cycles. Preservative-Free.
Shipping:	Shipping with blue ice.

### Antigen Details

Immunogen: Recombinant Protein: SARS-CoV Nucleoprotein / NP Protein (TMPY-02870)

Antigen Species: SARS

---

### Research Background

Coronaviruses are enveloped viruses with a positive-sense RNA genome and with a nucleocapsid of helical symmetry. Coronavirus nucleoproteins localize to the cytoplasm and the nucleolus, a subnuclear structure, in both virus-infected primary cells and in cells transfected with plasmids that express N protein. The coronavirus N protein is required for coronavirus RNA synthesis and has RNA chaperone activity that may be involved in template switch. Nucleocapsid protein is the most abundant protein of coronavirus. During virion assembly, N protein binds to viral RNA and leads to the formation of the helical nucleocapsid. Nucleocapsid protein is a highly immunogenic phosphoprotein also implicated in viral genome replication and in modulating cell signaling pathways. Because of the conservation of the N protein sequence and its strong immunogenicity, the N protein of coronavirus is chosen as a diagnostic tool.

**Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins**

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E\_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481