

Anti-Cortisol Binding Globulin Antibody (81857)

Product Details

| | |
|---------------|--------------|
| Ig Type: | Rabbit IgG |
| Reactivity: | Mouse |
| Conjugation: | Unconjugated |
| Clone: | 81857 |
| Purification: | Protein A |

Applications

| | |
|--------------|-----------------------|
| Application: | ELISA |
| Recommended | ELISA: 1:5000-1:10000 |

Properties

| | |
|----------------------|--|
| Stability & Storage: | Store at 2°C-8°C for 1 month. Store at -20°C or -80°C for 12 months. Avoid repeated freeze-thaw cycles. Preservative-Free. |
| Shipping: | Shipping with blue ice. |

Antigen Details

| | |
|------------------|---|
| Immunogen: | Recombinant Protein: Mouse SerpinA6 protein (TMPY-02107) |
| Antigen Species: | Mouse |
| Synonyms: | SerpinA6;serpin peptidase inhibitor, clade A (alpha-1 antiproteinase, antitrypsin), member 6; AI265318; Cbg; AV104445; serpin peptidase inhibitor, clade A (α -1 antiproteinase, antitrypsin), member 6 |

Research Background

Corticosteroid-binding globulin (CBG), also known as SerpinA6, is a non-inhibitory member of the serine proteinase inhibitor (serpin) superfamily. It is the high-affinity transport protein for glucocorticoids in vertebrate blood. CBG is specifically cleaved by this protease at a precise site close to its carboxy-terminus. This induces a conformation change and disrupts the binding between glucocorticoids and CBG, and promotes a significant and local release of glucocorticoids (over 90% of them are bound to CBG in human plasma). In this context, CBG directs glucocorticoids to sites of inflammation, and plays in consequence a crucial role in efficient glucocorticoid action in physiology. The SerpinA6 protein is mainly secreted by the liver. This negative acute phase protein regulates free cortisol levels in the blood and distributes cortisol to its target tissues. SerpinA6 deficiency is an extremely rare hereditary disorder characterized by reduced corticosteroid-binding capacity with normal or low plasma corticosteroid-binding globulin concentration, and normal or low basal cortisol levels associated with hypo-/hypertension and muscle fatigue. There are three heritable, human CBG gene mutations that can reduce CBG-cortisol binding affinity and/or reduce circulating CBG levels.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel: 781-999-4286 E_mail: info@targetmol.com Address: 34 Washington Street, Wellesley Hills, MA 02481