

Anti-CD16/FCGR3 Antibody-PE (40520)

Product Details

Ig Type:	Rabbit IgG
Reactivity:	Mouse
Conjugation:	PE
Clone:	40520
Purification:	Protein A

Applications

Verified Activity:	Flow cytometric analysis of Mouse FCGR3(CD16) expression on mouse splenocytes. BABL/c splenocytes were stained with PE-conjugated anti-Mouse FCGR3(CD16) and FITC-conjugated anti-Mouse CD3 (BD Pharmingen™ Cat. No. 553061). The dot plots were derived from events with the forward and side light-scatter characteristics of intact cells.
Application:	FCM
Recommended	10 µl/Test, 0.1 mg/ml

Properties

Stability & Storage:	Store at 2°C-8°C for 12 months, do not freeze. Keep away from direct sunlight.
Shipping:	Shipping with blue ice.

Antigen Details

Immunogen:	Recombinant Protein: Mouse CD16 / FCGR3 protein (TMPY-01834)
Antigen Species:	Mouse
Synonyms:	CD16;Fc receptor, IgG, low affinity III
Biology Area:	Fc Receptors

Research Background

Fc receptors bind the most common class of antibody, IgG, are called Fc gamma receptors (FcγR). FcγR is divided into three classes, Fc γ RI (CD64), Fc γ RII (CD32), and Fc γ RIII (CD16). CD16 protein is a multifunctional, low/intermediate affinity receptor, which belongs to the immunoglobulin superfamily. It is found on the surface of natural killer cells, neutrophil polymorphonuclear leukocytes, monocytes and macrophages. Mouse CD16 is encoded by a single gene, while, human CD16 is expressed as two distinct forms (CD16a/FcγRIIIa and CD16b/FcγRIIIb) encoded by two different highly homologous genes in a cell type-specific manner. CD16 is involved in phagocytosis, secretion of enzymes, inflammatory mediators, antibody-dependent cellular cytotoxicity (ADCC), and clearance of immune complexes.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only· Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481