

Anti-Endoglin/CD105 Antibody-PE (4U724)

Product Details

Ig Type:	Rabbit IgG
Reactivity:	Mouse
Conjugation:	PE
Clone:	4U724
Purification:	Protein A

Applications

Verified Activity:	Flow cytometric analysis of Mouse ENG(CD105) expression on bEnd.3 cells. Cells were stained with PE-conjugated anti-Mouse ENG(CD105). The fluorescence histograms were derived from gated events with the forward and side light-scatter characteristics of intact cells.
Application:	FCM
Recommended	5 µl/Test, 0.1 mg/ml

Properties

Stability & Storage:	Store at 2°C-8°C for 12 months, do not freeze. Keep away from direct sunlight. Sodium azide is toxic to cells and should be disposed of properly. Flush with large volumes of water during disposal.
Shipping:	Shipping with blue ice.

Antigen Details

Immunogen:	Recombinant Protein: Mouse Endoglin / CD105 / ENG Protein (TMPY-01615)
Antigen Species:	Mouse
Synonyms:	CD105;S-endoglin;AI528660;AI662476;Endo;endoglin
Biology Area:	Hemangioblast Markers, Cancer Drug Targets

Research Background

Endoglin, also known as CD105, is a type I homodimeric transmembrane glycoprotein with a large, disulfide-linked, extracellular region and a short, constitutively phosphorylated cytoplasmic tail. Endoglin contains an RGD tripeptide which is a key recognition structure in cellular adhesion, suggesting a critical role for endoglin in the binding of endothelial cells to integrins and/or other RGD receptors. Endoglin is highly expressed on vascular endothelial cells, chondrocytes, and syncytiotrophoblasts of term placenta. It is also found on activated monocytes, mesenchymal stem cells and leukemic cells of lymphoid and myeloid lineages. As an accessory receptor for the TGF- β superfamily ligands, endoglin binds TGF- β 1 and TGF- β 3 with high affinity not by itself but by associating with TGF- β type II receptor (T β RII) and activates the downstream signal pathways. In addition, in human umbilical vein endothelial cells, ALK-1 is also a receptor kinase for endoglin threonine phosphorylation, and mutations in either of the two genes result in the autosomal-dominant vascular dysplasia, hereditary hemorrhagic telangiectasia (HHT). Endoglin has been regarded as a powerful biomarker of neovascularization, and is associated with several solid tumor types.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481