

Anti-UCHL1 Antibody (6Z794)

Product Details

Ig Type:	Rabbit IgG
Reactivity:	Mouse
Conjugation:	Unconjugated
Clone:	6Z794
Purification:	Protein A

Applications

Verified Activity:	<p>1. Anti-UCHL1 rabbit monoclonal antibody at 1:500 dilution.</p> <ul style="list-style-type: none">-Lane A: 293T Whole Cell lysate.-Lysates/proteins at 30 µg per lane.-Secondary-Goat Anti-Rabbit IgG H&L (Dylight800) at 1/10000 dilution.-Developed using the Odyssey technique.-Performed under reducing conditions.-Predicted band size:25 kDa.-Observed band size:27 kDa. <p>2. Mouse UCHL1 was immunoprecipitated using:</p> <ul style="list-style-type: none">-Lane A:0.5 mg 293T Whole Cell Lysate.-Lane B:0.5 mg U87MG Whole Cell Lysate.-2 µL anti-Mouse UCHL1 rabbit monoclonal antibody and 15 µl of 50 % Protein G agarose.-Primary antibody:-Anti-Mouse UCHL1 rabbit monoclonal antibody, at 1:200 dilution.-Secondary antibody:-Dylight 800-labeled antibody to rabbit IgG (H+L), at 1:5000 dilution.-Developed using the odyssey technique.-Performed under reducing conditions.-Predicted band size: 25 kDa.-Observed band size: 25 kDa
Application:	IP,WB
Recommended	WB: 1:500-1:2000; IP: 0.5-2 µL/mg of lysate

Properties

Stability & Storage:	Store at 2°C-8°C for 1 month. Store at -20°C or -80°C for 12 months. Avoid repeated freeze-thaw cycles. Preservative-Free.
Shipping:	Shipping with blue ice.

Antigen Details

Immunogen: Recombinant Protein: Mouse UCHL1 protein (TMPY-02175)
Antigen Species: Mouse
Synonyms: ubiquitin carboxyl-terminal esterase L1 (ubiquitin thiolesterase)

Research Background

Ubiquitin carboxyl-terminal hydrolase isozyme L1, also known as UCH-L1, Ubiquitin thioesterase L1, PGP9.5 and UCHL1, is a deubiquitinating enzyme with important functions in recycling of ubiquitin. Regulated proteolysis by the ubiquitin pathway has been implicated in control of the cell cycle, transcriptional activation, cell fate and growth, and synaptogenesis. The ubiquitin-proteasome system is involved in synaptic plasticity and is proposed to be part of a molecular switch that converts short-term synaptic potentiation to long-term changes in synaptic strength. UCHL1 is found in neuronal cell bodies and processes throughout the neocortex (at protein level). It is expressed in neurons and cells of the diffuse neuroendocrine system and their tumors. UCHL1 is weakly expressed in ovary. UCHL1 is a ubiquitin-protein hydrolase. It is involved both in the processing of ubiquitin precursors and of ubiquitinated proteins. This enzyme is a thiol protease that recognizes and hydrolyzes a peptide bond at the C-terminal glycine of ubiquitin. UCHL1 also binds to free monoubiquitin and may prevent its degradation in lysosomes. The homodimer of UCHL1 may have ATP-independent ubiquitin ligase activity. UCHL1 dysfunction has been associated with neurodegeneration in Parkinson's, Alzheimer's, and Huntington's disease patients. Reduced UCHL1 function may jeopardize the survival of CNS neurons.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481