

Anti-UCHL3 Antibody (9J995)

Product Details

Ig Type:	Rabbit IgG
Reactivity:	Mouse
Conjugation:	Unconjugated
Clone:	9J995
Purification:	Protein A

Applications

	<p>1. Anti-UCHL3 rabbit monoclonal antibody at 1:500 dilution.</p> <ul style="list-style-type: none">-Lane A: Jurkat Whole Cell Lysate.-Lane B: K562 Whole Cell Lysate.-Lane C: 293T Whole Cell lysate.-Lysates/proteins at 30 µg per lane.-Secondary-Goat Anti-Rabbit IgG H&L (Dylight800) at 1/10000 dilution.-Developed using the Odyssey technique.-Performed under reducing conditions.-Predicted band size:26 kDa.-Observed band size:26 kDa(We are unsure as to the identity of these extra bands.)
Verified Activity:	<p>2. Mouse UCHL3 was immunoprecipitated using:</p> <ul style="list-style-type: none">-Lane A:0.5 mg K562 Whole Cell Lysate.-Lane B:0.5 mg 293T Whole Cell Lysate.-2 µL anti-Mouse UCHL3 rabbit monoclonal antibody and 15 µL of 50 % Protein G agarose.-Primary antibody:-Anti-Mouse UCHL3 rabbit monoclonal antibody, at 1:100 dilution.-Secondary antibody:-Dylight 800-labeled antibody to rabbit IgG (H+L), at 1:5000 dilution.-Developed using the odyssey technique.-Performed under reducing conditions.-Predicted band size: 26 kDa.-Observed band size: 26 kDa
Application:	IP,WB
Recommended	WB: 1:500-1:2000; IP: 0.5-2 µL/mg of lysate

Properties

Stability & Storage:	Store at 2°C-8°C for 1 month. Store at -20°C or -80°C for 12 months. Avoid repeated freeze-thaw cycles. Preservative-Free.
Shipping:	Shipping with blue ice.

Antigen Details

Immunogen: Recombinant Protein: Mouse UCHL3 Protein (TMPY-02190)
Antigen Species: Mouse
Synonyms: ubiquitin carboxyl-terminal esterase L3 (ubiquitin thiolesterase)

Research Background

Ubiquitin carboxyl-terminal hydrolase isozyme L3, also known as UCH-L3, Ubiquitin thioesterase L3 and UCHL3, is a ubiquitin-protein hydrolase that belongs to the peptidase C12 family. It is involved both in the processing of ubiquitin precursors and of ubiquitinated proteins. This enzyme is a thiol protease that recognizes and hydrolyzes a peptide bond at the C-terminal glycine of either ubiquitin or NEDD8. UCHL3 is highly expressed in heart, skeletal muscle, and testis. UCHL1 and UCHL3 are two of the deubiquitinating enzymes expressed in the brain. These phenotypes indicate the importance of UCHL1 and UCHL3 in the regulation of the central nervous system. UCHL3 functions as a de-ubiquitinating enzyme where lack of its hydrolase activity may result in the prominent accumulation of ubiquitinated proteins and subsequent induction of stress responses in skeletal muscle. UCHL3 has also been identified as a tumor-specific antigen in colon cancer.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel: 781-999-4286 E_mail: info@targetmol.com Address: 34 Washington Street, Wellesley Hills, MA 02481