

## Anti-CD6 Antibody-HRP (7C232)

## Product Details

|               |            |
|---------------|------------|
| Ig Type:      | Rabbit IgG |
| Reactivity:   | Mouse      |
| Conjugation:  | HRP        |
| Clone:        | 7C232      |
| Purification: | Protein A  |

## Applications

|              |                    |
|--------------|--------------------|
| Application: | ELISA              |
| Recommended  | ELISA: 0.1-1 µg/ml |

## Properties

|                      |  |
|----------------------|--|
| Stability & Storage: | Store at 2°C-8°C for 1 month. Store at -20°C or -80°C for 12 months. Avoid repeated freeze-thaw cycles. Preservative-Free. Keep away from direct sunlight. |
| Shipping:            | Shipping with blue ice.  |

## Antigen Details

|                  |  |
|------------------|--|
| Immunogen:       | Recombinant Protein: Mouse CD6 / Cluster of Differentiation 6 Protein (TMPY-02254) |
| Antigen Species: | Mouse  |
| Synonyms:        | CD6 molecule;TP120   |
| Biology Area:    | ITIM/ITAM Immunoreceptors and Related Molecules                                    |

## Research Background

T-cell differentiation antigen CD6, also known as TP12 and CD6, is a single-pass type I membrane protein which contains three SRCR domains. CD6 / TP12 is a cell surface glycoprotein expressed primarily on T cells, it may function as a costimulatory molecule and may play a role in autoreactive immune responses. CD6 / TP12 is expressed by thymocytes, mature T-cells, a subset of B-cells known as B-1 cells, and by some cells in the brain. CD6 ligand termed CD166 (previously known as activated leukocyte cell adhesion molecule, ALCAM) has been identified and shown to be expressed on activated T cells, B cells, thymic epithelium, keratinocytes, and in rheumatoid arthritis synovial tissue. CD6 / TP12 binds to activated leukocyte cell adhesion molecule (CD166), and is considered as a costimulatory molecule involved in lymphocyte activation and thymocyte development. CD6 / TP12 partially associates with the TCR / CD3 complex and colocalizes with it at the center of the mature immunological synapse (IS) on T lymphocytes. During thymic development CD6-dependent signals may contribute both to thymocyte survival, and to the overall functional avidity of selection in both man and mouse.

**Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins**

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel: 781-999-4286 E\_mail: info@targetmol.com Address: 34 Washington Street, Wellesley Hills, MA 02481