

Anti-CD6 Antibody-FITC (9H618)

Product Details

| | |
|---------------|------------|
| Ig Type: | Rabbit IgG |
| Reactivity: | Mouse |
| Conjugation: | FITC |
| Clone: | 9H618 |
| Purification: | Protein A |

Applications

| | |
|--------------------|---------------------------------------------------|
| Verified Activity: | Analysis of CD6 expression on spleen lymphocytes. |
| Application: | FCM |
| Recommended | 2 µl/Test, 0.1 mg/ml |

Properties

| | |
|----------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Stability & Storage: | Store at 2°C-8°C for 12 months, do not freeze. Keep away from direct sunlight. Sodium azide is toxic to cells and should be disposed of properly. Flush with large volumes of water during disposal. |
| Shipping: | Shipping with blue ice. |

Antigen Details

| | |
|------------------|-----------------------------------------------------|
| Immunogen: | Recombinant Protein: Mouse CD6 protein (TMPY-02254) |
| Antigen Species: | Mouse |
| Synonyms: | TP120;CD6 molecule |
| Biology Area: | ITIM/ITAM Immunoreceptors and Related Molecules |

Research Background

T-cell differentiation antigen CD6, also known as TP12 and CD6, is a single-pass type I membrane protein which contains three SRCR domains. CD6 / TP12 is a cell surface glycoprotein expressed primarily on T cells, it may function as a costimulatory molecule and may play a role in autoreactive immune responses. CD6 / TP12 is expressed by thymocytes, mature T-cells, a subset of B-cells known as B-1 cells, and by some cells in the brain. CD6 ligand termed CD166 (previously known as activated leukocyte cell adhesion molecule, ALCAM) has been identified and shown to be expressed on activated T cells, B cells, thymic epithelium, keratinocytes, and in rheumatoid arthritis synovial tissue. CD6 / TP12 binds to activated leukocyte cell adhesion molecule (CD166), and is considered as a costimulatory molecule involved in lymphocyte activation and thymocyte development. CD6 / TP12 partially associates with the TCR / CD3 complex and colocalizes with it at the center of the mature immunological synapse (IS) on T lymphocytes. During thymic development CD6-dependent signals may contribute both to thymocyte survival, and to the overall functional avidity of selection in both man and mouse.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481