

## Anti-IL-1R1 Antibody-APC (3F72)

## Product Details

Ig Type:	Rabbit IgG
Reactivity:	Mouse
Conjugation:	APC
Clone:	3F72
Purification:	Protein A

## Applications

Verified Activity:	Flow cytometric analysis of Mouse IL1R1(CD121a) expression on NIH/3T3 cells. NIH/3T3 cells were stained with APC-conjugated anti-mouse IL1R1(CD121a). The histogram were derived from gated events with the forward and side light-scatter characteristics of intact cells.
Application:	FCM
Recommended	10 µl/Test, 0.1 mg/ml

## Properties

Stability & Storage:	Store at 2°C-8°C for 12 months, do not freeze. Keep away from direct sunlight.
Shipping:	Shipping with blue ice.

## Antigen Details

Immunogen:	Recombinant Protein: Mouse IL-1R1 / CD121a protein (TMPY-02341)
Antigen Species:	Mouse
Synonyms:	interleukin 1 receptor, type I
Biology Area:	Neuroinflammation

## Research Background

Interleukin 1 receptor, type I (IL-1R1) also known as CD121a (Cluster of Differentiation 121a), is an interleukin receptor. IL-1R1/CD121a is a cytokine receptor that belongs to the interleukin 1 receptor family. This protein is a receptor for interleukin alpha (IL1A), interleukin beta (IL1B), and interleukin 1 receptor, type I (IL1R1/IL1RA). IL-1R1/CD121a is an important mediator involved in many cytokines induced immune and inflammatory responses. This protein has been characterized by pharmacological and molecular techniques in the mouse brain. The spindle-shaped astrocytes enclose the wound, separating the healthy from damaged neural tissue. The shape change and subsequent repair processes are IL-1 $\beta$  activity-dependent, acting through the IL-1 type 1 receptor (IL-1R1), as co-application of the IL-1 type 1 receptor antagonist protein (IL-1ra) blocks IL-1 $\beta$  induced effects. In the spleen, a slight increase in IL-1R ACP and IL-1R1 was observed during the first hours following LPS stimulation. In conclusion, IL-1R ACP mRNA is expressed in the brain and in other tissues where IL-1R1/CD121a transcripts are found. However, the regulation of its expression is distinct from IL-1R1/CD121a. The high level of expression and the lack of regulation of IL-1R ACP transcripts in the brain under inflammatory conditions suggest that the protein might be constitutively expressed in excess.

**Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins**

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E\_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481