

Anti-SIRP alpha Antibody-APC (2E26)

Product Details

Ig Type:	Rabbit IgG
Reactivity:	Mouse
Conjugation:	APC
Clone:	2E26
Purification:	Protein A

Applications

Verified Activity:	Flow cytometric analysis of Mouse SIRPA(CD172a) expression on BABL/c bone marrow cells. Cells were stained with APC-conjugated anti-Mouse SIRPA(CD172a). The fluorescence histograms were derived from gated events with the forward and side light-scatter characteristics of intact cells.
Application:	FCM
Recommended	5 µl/Test, 0.1 mg/ml

Properties

Stability & Storage:	Store at 2°C-8°C for 12 months, do not freeze. Keep away from direct sunlight.
Shipping:	Shipping with blue ice.

Antigen Details

Immunogen:	Recombinant Protein: Mouse SIRP alpha / CD172a Protein (TMPY-02632)
Antigen Species:	Mouse
Synonyms:	SHPS1;P84;SIRP α;CD172A;MYD-1;signal-regulatory protein α;MFR;PTPNS1;signal-regulatory protein alpha;SIRP alpha;SIRP;BIT
Biology Area:	Cancer Drug Targets

Research Background

Tyrosine-protein phosphatase non-receptor type substrate 1, also known as SHP substrate 1, Inhibitory receptor SHPS-1, Brain Ig-like molecule with tyrosine-based activation motifs, Macrophage fusion receptor, CD172 antigen-like family member A, SIRPA and CD172a, is a single-pass type I membrane protein which contains two Ig-like C1-type (immunoglobulin-like) domains and one Ig-like V-type (immunoglobulin-like) domain. SIRPA is ubiquitously expressed. It is highly expressed in brain and detected at lower levels in heart, placenta, lung, testis, ovary, colon, liver, small intestine, prostate, spleen, kidney, skeletal muscle and pancreas. It is also detected on myeloid cells, but not T-cells. SIRPA is an immunoglobulin-like cell surface receptor for CD47. SIRPA acts as docking protein and induces translocation of PTPN6, PTPN11 and other binding partners from the cytosol to the plasma membrane. SIRPA supports adhesion of cerebellar neurons, neurite outgrowth and glial cell attachment. It may play a key role in intracellular signaling during synaptogenesis and in synaptic function. SIRPA is involved in the negative regulation of receptor tyrosine kinase-coupled cellular responses induced by cell adhesion, growth factors or insulin. It mediates negative regulation of phagocytosis, mast cell activation and dendritic cell activation. Cancer Immunotherapy Co-inhibitory Immune Checkpoint Targets Immune Checkpoint Immune Checkpoint Detection: ELISA Antibodies Immune Checkpoint Detection: FCM Antibodies Immune Checkpoint Detection: IHC Antibodies Immune

A DRUG SCREENING EXPERT

Checkpoint Detection: IP AntibodiesImmune Checkpoint Detection: WB AntibodiesImmune Checkpoint
TargetsImmunotherapyTargeted Therapy

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only· Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481