

## Anti-NGAL/Lipocalin-2 Antibody-HRP (40488)

## Product Details

Ig Type:	Mouse IgG2a
Reactivity:	Rat
Conjugation:	HRP
Clone:	40488
Purification:	Protein A

## Applications

Application:	ELISA
Recommended	ELISA: 0.1-1 µg/mL

## Properties

Stability & Storage:	Store at 2°C-8°C for 1 month. Store at -20°C or -80°C for 12 months. Avoid repeated freeze-thaw cycles. Preservative-Free. Keep away from direct sunlight.
Shipping:	Shipping with blue ice.

## Antigen Details

Immunogen:	Recombinant Protein: Rat NGAL / LCN2 / Lipocalin-2 Protein (TMPY-02316)
Antigen Species:	Rat
Synonyms:	NGAL;Lipocalin-2;p25;Oncogene 24p3;MSFI;Siderocalin LCN2

## Research Background

Lipocalin-2 (LCN2), also known as neutrophil gelatinase-associated lipocalin (NGAL), is a 25 kDa protein belonging to the lipocalin superfamily. It was initially found in activated neutrophils, however, many other cells, like kidney tubular cells, may produce NGAL in response to various insults. This protein is released from injured tubular cells after various damaging stimuli, is already known by nephrologists as one of the most promising biomarkers of incoming Acute Kidney Injury (AKI). Recent evidence also suggests its role as a biomarker in a variety of other renal and non-renal conditions. Moreover, recent studies seem to suggest a potential involvement of this factor also in the genesis and progression of chronic kidney diseases. NGAL is the first known mammalian protein that specifically binds organic molecules called siderophores, which are high-affinity iron chelators. NGAL, first known as an antibacterial factor of natural immunity, and an acute-phase protein, is currently one of the most interesting and enigmatic proteins involved in the process of tumor development. acting as an intracellular iron carrier and protecting MMP9 from proteolytic degradation, NGAL has a clear pro-tumoral effect, as has already been observed in different tumors (e.g. breast, stomach, esophagus, brain) in humans. In thyroid carcinomas, NGAL is strongly induced by NF-κB, an important factor involved both in tumor growth and in the link between chronic inflammation and neoplastic development. Thus, Lipocalin-2 (LCN2/NGAL) has been implicated in a variety of processes including cell differentiation, proliferation, survival, and morphogenesis.

**Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins**

This product is for Research Use Only. Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E\_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481