

Anti-BCAM Antibody (5W424)

Product Details

Ig Type:	Rabbit IgG
Reactivity:	Rat
Conjugation:	Unconjugated
Clone:	5W424
Purification:	Protein A

Applications

Verified Activity:	1. Immunochemical staining of rat BCAM in rat skin with rabbit monoclonal antibody (1:200, formalin-fixed paraffin embedded sections).
	2. Immunochemical staining of rat BCAM in rat liver with rabbit monoclonal antibody (1:200, formalin-fixed paraffin embedded sections).
	3. Immunochemical staining of rat BCAM in rat kidney with rabbit monoclonal antibody (1:200, formalin-fixed paraffin embedded sections).
Application:	IHC-P
Recommended	IHC-P: 1:100-1:500

Properties

Stability & Storage:	Store at 2°C-8°C for 1 month. Store at -20°C or -80°C for 12 months. Avoid repeated freeze-thaw cycles. Preservative-Free.
Shipping:	Shipping with blue ice.

Antigen Details

Immunogen:	Recombinant Protein: Rat BCAM / CD239 protein (TMPY-02999)
Antigen Species:	Rat
Synonyms:	basal cell adhesion molecule (Lutheran blood group)

Research Background

The Lutheran (Lu) blood group and basal cell adhesion molecule (BCAM) antigens are both carried by 2 glycoprotein isoforms of the immunoglobulin superfamily representing receptors for the laminin alpha(5) chain. It is a transmembrane receptor with five immunoglobulin-like domains in its extracellular region, and is therefore classified as a member of the immunoglobulin (Ig) gene family. In addition to red blood cells, Lu/BCAM proteins are expressed in endothelial cells of vascular capillaries and in epithelial cells of several tissues. BCAM/LU has a wide tissue distribution with a predominant expression in the basal layer of the epithelium and the endothelium of blood vessel walls. As designated as CD239 recently, BCAM and LU share a significant sequence similarity with the CD146 (MUC18) and CD166, and themselves are adhesion molecules that bind laminin with high affinity. Laminins are found in all basement membranes and are involved in cell differentiation, adhesion, migration, and proliferation. BCAM is upregulated following malignant transformation of some cell types in vivo and in vitro, thus being a candidate molecule involved in tumor progression. In addition, BCAM interacts with integrin in sickle red cells, and thus may potentially play a role in vaso-occlusive episodes.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481