

## Anti-COMT Antibody (6P98)

## Product Details

Ig Type:	Rabbit IgG
Reactivity:	Human
Conjugation:	Unconjugated
Clone:	6P98
Purification:	Affinity-chromatography

## Applications

Verified Activity:	1. Western Blot -Positive WB detected in: 293 whole cell lysate, THP-1 whole cell lysate, HepG2 whole cell lysate -All lanes: COMT Antibody at 1:1000 -Secondary: Goat polyclonal to rabbit IgG at 1/50000 dilution -Predicted band size: 31, 25 kDa -Observed band size: 31 kDa
	2. IHC image of TMAH-00292 diluted at 1:100 and staining in paraffin-embedded human brain tissue performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a Goat anti-rabbit IgG polymer labeled by HRP and visualized using 0.05% DAB.
Application:	ELISA, WB, IHC
Recommended	WB:1:500-1:5000; IHC:1:50-1:200.

## Properties

Stability & Storage:	Store at -20°C or -80°C for 12 months. Avoid repeated freeze-thaw cycles.
Shipping:	Shipping with blue ice.

## Antigen Details

Immunogen:	A synthetic peptide: Human COMT
Antigen Species:	Human
Gene ID:	1312
Uniprot ID:	P21964
Synonyms:	HEL-S-98n;catechol-O-methyltransferase
Biology Area:	Neuroscience, Cancer, Metabolism, Signal transduction

## Research Background

Catalyzes the O-methylation, and thereby the inactivation, of catecholamine neurotransmitters and catechol hormones. Also shortens the biological half-lives of certain neuroactive drugs, like L-DOPA, alpha-methyl DOPA and isoproterenol.

**Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins**

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E\_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481