

Anti-GDF-15 Antibody (2W692)

Product Details

Ig Type:	Rabbit IgG
Reactivity:	Human
Conjugation:	Unconjugated
Clone:	2W692
Purification:	Affinity-chromatography

Applications

Verified Activity:	IHC image of TMAH-00481 diluted at 1:100 and staining in paraffin-embedded human placenta tissue performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a Goat anti-rabbit IgG polymer labeled by HRP and visualized using 0.05% DAB.
Application:	ELISA,IHC
Recommended	IHC:1:50-1:200.

Properties

Stability & Storage:	Store at -20°C or -80°C for 12 months. Avoid repeated freeze-thaw cycles.
Shipping:	Shipping with blue ice.

Antigen Details

Immunogen:	A synthetic peptide: Human GDF15
Antigen Species:	Human
Gene ID:	9518
Uniprot ID:	Q99988
Synonyms:	MIC-1;PTGFB;PTGF-beta;Placental TGF-beta;MIC1;Placental TGF-β;RG-1;PTGF-β;PLAB;NAG-1;PDF;PTGFBPTGF-β;PTGFBPTGF-beta;GDF-15;GDF15
Biology Area:	Cardiovascular, Cell biology, Signal transduction

Research Background

Regulates food intake, energy expenditure and body weight in response to metabolic and toxin-induced stresses. Binds to its receptor, GFRAL, and activates GFRAL-expressing neurons localized in the area postrema and nucleus tractus solitarius of the brainstem. It then triggers the activation of neurons localized within the parabrachial nucleus and central amygdala, which constitutes part of the 'emergency circuit' that shapes feeding responses to stressful conditions. On hepatocytes, inhibits growth hormone signaling.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only. Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481