

Anti-IL-1 beta/IL-1F2 Antibody (4Y981)

Product Details

Ig Type:	hIgG1
Reactivity:	Human
Conjugation:	Unconjugated
Clone:	4Y981
Purification:	Affinity-chromatography

Applications

1. IHC image of TMAH-00616 diluted at 1:100 and staining in paraffin-embedded human kidney tissue performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a Goat anti-Human IgG labeled by HRP and visualized using 0.05% DAB.

Verified Activity: 2. Overlay Peak curve showing PC3 cells stained with TMAH-00616 (red line) at 1:200. The cells were fixed in 4% formaldehyde and permeated by 0.2% TritonX-100. Then 10% normal goat serum to block non-specific protein-protein interactions followed by the antibody (1 μ g/1*10⁶ cells) for 45min at 4°C. The secondary antibody used was FITC-conjugated Goat Anti-human IgG (H+L) at 1:200 dilution for 35min at 4°C. Control antibody (green line) was human IgG (1 μ g/1*10⁶ cells) used under the same conditions. Acquisition of >10,000 events was performed.

Application: ELISA,FCM,IHC

Recommended IHC:1:50-1:200; FCM:1:50-1:200.

Properties

Stability & Storage: Store at -20°C or -80°C for 12 months. Avoid repeated freeze-thaw cycles.

Shipping: Shipping with blue ice.

Antigen Details

Immunogen:	Recombinant Protein: Human IL1B Protein
Antigen Species:	Human
Gene ID:	3553
Uniprot ID:	P01584
Synonyms:	IL1F2;IL-1BETA;IL-1 β ;IL1B;IL-1 β /IL-1F2;IL-1 beta;Interleukin-1 β ;IL-1 β ;Interleukin-1 beta;IL-1B
Biology Area:	Cardiovascular, Immunology, Metabolism

Research Background

Potent proinflammatory cytokine. Initially discovered as the major endogenous pyrogen, induces prostaglandin synthesis, neutrophil influx and activation, T-cell activation and cytokine production, B-cell activation and antibody

A DRUG SCREENING EXPERT

production, and fibroblast proliferation and collagen production. Promotes Th17 differentiation of T-cells. Synergizes with IL12/interleukin-12 to induce IFNG synthesis from T-helper 1 (Th1) cells. Plays a role in angiogenesis by inducing VEGF production synergistically with TNF and IL6.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481