

Anti-Phospho-Histone H3.1(Ser10) Antibody (6R499)

Product Details

Ig Type:	Rabbit IgG
Reactivity:	Human
Conjugation:	Unconjugated
Clone:	6R499
Purification:	Affinity-chromatography

Applications

Verified Activity:	<p>1. Western Blot</p> <ul style="list-style-type: none">-Positive WB detected in Hela whole cell lysate, MCF-7 whole cell lysate, PC3 whole cell lysate-All lanes Phospho-Histone H3.1(S10)antibody at 0.5µg/ml-Secondary: Goat polyclonal to rabbit IgG at 1/50000 dilution-Predicted band size: 15 KDa-Observed band size: 15 KDa <p>2. Immunocytochemistry analysis of TMAH-00907 diluted at 1:100 and staining in Hela cells performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.</p> <p>3. Immunofluorescence staining of Hela cells with TMAH-00907 at 1:31, counter-stained with DAPI. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4°C. The secondary antibody was Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG (H+L).</p>
Application:	ELISA, ICC, IF, WB
Recommended	WB: 1:500-1:2000; ICC: 1:50-1:500; IF: 1:30-1:200.

Properties

Stability & Storage:	Store at -20°C or -80°C for 12 months. Avoid repeated freeze-thaw cycles.
Shipping:	Shipping with blue ice.

Antigen Details

Immunogen:	A synthetic peptide
Antigen Species:	Human
Gene ID:	8350
Uniprot ID:	P68431
Synonyms:	p-Histone H3.1(S10);histone cluster 1, H3a;Histone H3.1(p-S10);Histone H3;Histone H3.1(p-Ser10);p-Histone H3.1(Ser10);Histone H3.1;Phospho-Histone H3.1(S10);HIST1H3A, HIST1H3B, HIST1H3C, HIST1H3D, HIST1H3E, HIST1H3F, HIST1H3G, HIST1H3H, HIST1H3I, HIST1H3J
Biology Area:	Epigenetics and Nuclear Signaling

Research Background

Core component of nucleosome. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of histones, also called histone code, and nucleosome remodeling.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481