

Anti-PKN2 Antibody (7R455)

Product Details

Ig Type:	Rabbit IgG
Reactivity:	Human
Conjugation:	Unconjugated
Clone:	7R455
Purification:	Affinity-chromatography

Applications

	<p>1. Western Blot</p> <ul style="list-style-type: none">-Positive WB detected in: Hela whole cell lysate, 293T whole cell lysate-All lanes: EB13 antibody at 1:1000-Secondary: Goat polyclonal to rabbit IgG at 1/50000 dilution-Predicted band size: 140 kDa-Observed band size: 140 kDa
Verified Activity:	<p>2. Overlay Peak curve showing HepG2 cells stained with TMAH-00987 (red line) at 1:50. The cells were fixed in 4% formaldehyde and permeated by 0.2% TritonX-100. Then 10% normal goat serum to block non-specific protein-protein interactions followed by the antibody (1µg/1*10⁶ cells) for 45min at 4°C. The secondary antibody used was FITC-conjugated Goat Anti-rabbit IgG(H+L) at 1:200 dilution for 35min at 4°C. Control antibody (green line) was rabbit IgG (1µg/1*10⁶ cells) used under the same conditions. Acquisition of >10,000 events was performed.</p>
Application:	ELISA,FCM,WB
Recommended	WB:1:500-1:2000; FCM:1:50-1:200.

Properties

Stability & Storage:	Store at -20°C or -80°C for 12 months. Avoid repeated freeze-thaw cycles.
Shipping:	Shipping with blue ice.

Antigen Details

Immunogen:	A synthetic peptide: Human PKN2
Antigen Species:	Human
Gene ID:	5586
Uniprot ID:	Q16513
Synonyms:	PKN gamma;Serine/threonine-protein kinase N2;PRK2;Protein-kinase C-related kinase 2;PKN 2;EC 2.7.11.13;PRKCL2;Protein kinase C-like 2
Biology Area:	Signal transduction

Research Background

PKC-related serine/threonine-protein kinase and Rho/Rac effector protein that participates in specific signal transduction responses in the cell. Plays a role in the regulation of cell cycle progression, actin cytoskeleton

A DRUG SCREENING EXPERT

assembly, cell migration, cell adhesion, tumor cell invasion and transcription activation signaling processes. Phosphorylates CTTN in hyaluronan-induced astrocytes and hence decreases CTTN ability to associate with filamentous actin. Phosphorylates HDAC5, therefore lead to impair HDAC5 import. Direct RhoA target required for the regulation of the maturation of primordial junctions into apical junction formation in bronchial epithelial cells. Required for G2/M phases of the cell cycle progression and abscission during cytokinesis in a ECT2-dependent manner. Stimulates FYN kinase activity that is required for establishment of skin cell-cell adhesion during keratinocytes differentiation. Regulates epithelial bladder cells speed and direction of movement during cell migration and tumor cell invasion. Inhibits Akt pro-survival-induced kinase activity. Mediates Rho protein-induced transcriptional activation via the c-fos serum response factor (SRF). Involved in the negative regulation of ciliogenesis. (Microbial infection) Phosphorylates HCV NS5B leading to stimulation of HCV RNA replication.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481