

## Anti-PTBP1 Antibody (9D226)

### Product Details

Ig Type:	Rabbit IgG
Reactivity:	Human
Conjugation:	Unconjugated
Clone:	9D226
Purification:	Affinity-chromatography

### Applications

Verified Activity:	<p>1. Western Blot</p> <ul style="list-style-type: none"><li>-Positive WB detected in: Hela whole cell lysate, HepG2 whole cell lysate, PC-3 whole cell lysate, K562 whole cell lysate, HL-60 whole cell lysate, MCF-7 whole cell lysate, Jurkat whole cell lysate</li><li>-All lanes: PTBP1 antibody at 1:1000</li><li>-Secondary: Goat polyclonal to rabbit IgG at 1/50000 dilution</li><li>-Predicted band size: 58, 60, 60 kDa</li><li>-Observed band size: 58 kDa</li></ul> <p>2. IHC image of TMAH-01012 diluted at 1:100 and staining in paraffin-embedded human breast cancer performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a Goat anti-rabbit IgG polymer labeled by HRP and visualized using 0.05% DAB.</p> <p>3. IHC image of TMAH-01012 diluted at 1:100 and staining in paraffin-embedded human testis tissue performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a Goat anti-rabbit IgG polymer labeled by HRP and visualized using 0.05% DAB.</p>
Application:	ELISA,IHC,WB
Recommended	WB:1:500-1:5000; IHC:1:50-1:200.

### Properties

Stability & Storage:	Store at -20°C or -80°C for 12 months. Avoid repeated freeze-thaw cycles.
Shipping:	Shipping with blue ice.

### Antigen Details

Immunogen:	A synthetic peptide: Human PTBP1
Antigen Species:	Human
Gene ID:	5725
Uniprot ID:	P26599
Synonyms:	hnRNP I;PTBP 1;57 kDa RNA-binding protein PPTB-1;PTB;Polypyrimidine tract-binding protein 1;Heterogeneous nuclear ribonucleoprotein I
Biology Area:	Epigenetics and Nuclear Signaling

---

### Research Background

Plays a role in pre-mRNA splicing and in the regulation of alternative splicing events. Activates exon skipping of its own pre-mRNA during muscle cell differentiation. Binds to the polypyrimidine tract of introns. May promote RNA looping when bound to two separate polypyrimidine tracts in the same pre-mRNA. May promote the binding of U2 snRNP to pre-mRNA. Cooperates with RAVR1 to modulate switching between mutually exclusive exons during maturation of the TPM1 pre-mRNA. Represses the splicing of MAPT/Tau exon 10. In case of infection by picornaviruses, binds to the viral internal ribosome entry site (IRES) and stimulates the IRES-mediated translation.

**Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins**

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E\_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481