

Anti-S6K1/RPS6KB1 Antibody (9L715)

Product Details

Ig Type:	Rabbit IgG
Reactivity:	Human, Mouse
Conjugation:	Unconjugated
Clone:	9L715
Purification:	Affinity-chromatography

Applications

	Western Blot
	-Positive WB detected in: NIH/3T3 whole cell lysate, MCF-7 whole cell lysate, Hela whole cell lysate, 293T whole cell lysate, Mouse Brain whole cell lysate
Verified Activity:	-All lanes: S6K1 antibody at 1:1000 -Secondary: Goat polyclonal to rabbit IgG at 1/50000 dilution -Predicted band size: 60, 57, 53, 57, 52 kDa -Observed band size: 70 kDa
Application:	ELISA, WB
Recommended	WB:1:500-1:5000.

Properties

Stability & Storage:	Store at -20°C or -80°C for 12 months. Avoid repeated freeze-thaw cycles.
Shipping:	Shipping with blue ice.

Antigen Details

Immunogen:	A synthetic peptide: Human S6K1
Antigen Species:	Human
Gene ID:	6198
Uniprot ID:	P23443
Synonyms:	S6K1;p70(S6K)-alpha;p70(S6K)-α;PS6K;S6K;p70-α;STK14A;p70-alpha;p70-S6K;RPS6KB1;p70 S6KA;S6K-β-1;ribosomal protein S6 kinase, polypeptide 1;S6K-beta-1
Biology Area:	Epigenetics and Nuclear Signaling, Cell biology, Metabolism, Signal transduction

Research Background

Serine/threonine-protein kinase that acts downstream of mTOR signaling in response to growth factors and nutrients to promote cell proliferation, cell growth and cell cycle progression. Regulates protein synthesis through phosphorylation of EIF4B, RPS6 and EEF2K, and contributes to cell survival by repressing the pro-apoptotic function of BAD. Under conditions of nutrient depletion, the inactive form associates with the EIF3 translation initiation complex. Upon mitogenic stimulation, phosphorylation by the mammalian target of rapamycin complex 1 (mTORC1) leads to dissociation from the EIF3 complex and activation. The active form then phosphorylates and activates several substrates in the pre-initiation complex, including the EIF2B complex and the cap-binding complex component EIF4B. Also controls translation initiation by phosphorylating a negative regulator of EIF4A, PDCD4,

targeting it for ubiquitination and subsequent proteolysis. Promotes initiation of the pioneer round of protein synthesis by phosphorylating POLDIP3/SKAR. In response to IGF1, activates translation elongation by phosphorylating EEF2 kinase (EEF2K), which leads to its inhibition and thus activation of EEF2. Also plays a role in feedback regulation of mTORC2 by mTORC1 by phosphorylating RICTOR, resulting in the inhibition of mTORC2 and AKT1 signaling. Mediates cell survival by phosphorylating the pro-apoptotic protein BAD and suppressing its pro-apoptotic function. Phosphorylates mitochondrial URI1 leading to dissociation of a URI1-PPP1CC complex. The free mitochondrial PPP1CC can then dephosphorylate RPS6KB1 at Thr-412, which is proposed to be a negative feedback mechanism for the RPS6KB1 anti-apoptotic function. Mediates TNF-alpha-induced insulin resistance by phosphorylating IRS1 at multiple serine residues, resulting in accelerated degradation of IRS1. In cells lacking functional TSC1-2 complex, constitutively phosphorylates and inhibits GSK3B. May be involved in cytoskeletal rearrangement through binding to neurabin. Phosphorylates and activates the pyrimidine biosynthesis enzyme CAD, downstream of MTOR. Following activation by mTORC1, phosphorylates EPRS and thereby plays a key role in fatty acid uptake by adipocytes and also most probably in interferon-gamma-induced translation inhibition.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel: 781-999-4286 E_mail: info@targetmol.com Address: 34 Washington Street, Wellesley Hills, MA 02481