

Anti-TERT Antibody (3E772)

Product Details

| | |
|---------------|-------------------------|
| Ig Type: | Rabbit IgG |
| Reactivity: | Human |
| Conjugation: | Unconjugated |
| Clone: | 3E772 |
| Purification: | Affinity-chromatography |

Applications

| | |
|--------------------|--|
| Verified Activity: | IHC image of TMAH-01160 diluted at 1:100 and staining in paraffin-embedded human lung cancer performed on a Leica Bond TM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a Goat anti-rabbit IgG polymer labeled by HRP and visualized using 0.05% DAB. |
| Application: | ELISA, IHC |
| Recommended | IHC:1:50-1:200. |

Properties

| | |
|----------------------|---|
| Stability & Storage: | Store at -20°C or -80°C for 12 months. Avoid repeated freeze-thaw cycles. |
| Shipping: | Shipping with blue ice. |

Antigen Details

| | |
|------------------|--|
| Immunogen: | A synthetic peptide: Human TERT |
| Antigen Species: | Human |
| Gene ID: | 7015 |
| Uniprot ID: | O14746 |
| Biology Area: | Epigenetics and Nuclear Signaling, Cancer, Developmental biology, Stem cells |

Research Background

Telomerase is a ribonucleoprotein enzyme essential for the replication of chromosome termini in most eukaryotes. Active in progenitor and cancer cells. Inactive, or very low activity, in normal somatic cells. Catalytic component of the telomerase holoenzyme complex whose main activity is the elongation of telomeres by acting as a reverse transcriptase that adds simple sequence repeats to chromosome ends by copying a template sequence within the RNA component of the enzyme. Catalyzes the RNA-dependent extension of 3'-chromosomal termini with the 6-nucleotide telomeric repeat unit, 5'-TTAGGG-3'. The catalytic cycle involves primer binding, primer extension and release of product once the template boundary has been reached or nascent product translocation followed by further extension. More active on substrates containing 2 or 3 telomeric repeats. Telomerase activity is regulated by a number of factors including telomerase complex-associated proteins, chaperones and polypeptide modifiers. Modulates Wnt signaling. Plays important roles in aging and antiapoptosis.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481